

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>5</sup> : C07D 211/56, 417/12, 401/12 C07D 487/08, 453/02 A61K 31/445, 31/435 // (C07D 453/02, 221:00, 211:00) (C07D 487/08, 209:00, 209:00)</p>	A1	<p>(11) International Publication Number: <b>WO 94/04496</b></p> <p>(43) International Publication Date: 3 March 1994 (03.03.94)</p>
<p>(21) International Application Number: PCT/US93/04063</p> <p>(22) International Filing Date: 5 May 1993 (05.05.93)</p> <p>(30) Priority data: 07/932,392 19 August 1992 (19.08.92) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 07/932,392 (CIP) Filed on 19 August 1992 (19.08.92)</p> <p>(71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).</p>		<p>(72) Inventors; and (75) Inventors/Applicants (for US only) : HOWARD, Harry, R., Jr. [US/US]; 272 Westwoods Terrace, Bristol, CT 06010 (US). IKUNAKA, Masaya [JP/JP]; ITO, Fumitaka [JP/JP]; 1-1-2, Sakuragaoka, Taketoyo, Chita, Aichi 470-23 (JP). LOWE, John, A., III [US/US]; 28 Coveside Lane, Stonington, CT 06378 (US). NAKANE, Masami [JP/JP]; 6-28, Kawanahonmachi, Showa-ku, Nagoya, Aichi 466 (JP). O'NEILL, Brian, T. [US/US]; 1526 Essex Road, Westbrook, CT 06498 (US). ROSEN, Terry, R. [US/US]; 245 Grassy Hill Road, East Lyme, CT 06333 (US). SATAKE, Kunio [JP/JP]; 4-58, Hakusan, Handa, Aichi 475 (JP).</p> <p>(74) Agents: RICHARDSON, Peter, C. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).</p> <p>(81) Designated States: AU, BR, CA, CZ, JP, KR, NO, NZ, PL, RU, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published With international search report.</p>
<p>(54) Title: SUBSTITUTED BENZYLAMINO NITROGEN CONTAINING NON-AROMATIC HETEROCYCLES</p> <div style="text-align: center; margin: 20px 0;"> <p style="margin-left: 150px;">(I)</p> </div> <p>(57) Abstract</p> <p>The present invention relates to novel substituted benzylamino nitrogen containing non-aromatic heterocycles and, specifically, to compounds of formula (I) wherein W, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and A are as defined in the specification, and to intermediates used in the synthesis of such compounds. The novel compounds of formula (I) are useful in the treatment of inflammatory and central nervous system disorders, as well as other disorders.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

-1-

5                    SUBSTITUTED BENZYLAMINO NITROGEN CONTAINING  
                      NON-AROMATIC HETEROCYCLES

Background of the Invention

                      The present invention relates to novel substituted  
benzylamino nitrogen containing non-aromatic heterocycles,  
10    pharmaceutical compositions comprising such compounds and  
the use of such compounds in the treatment and prevention of  
inflammatory and central nervous system disorders, as well  
as several other disorders. The pharmaceutically active  
compounds of this invention are substance P receptor  
15    antagonists. This invention also relates to novel  
intermediates used in the synthesis of such substance P  
receptor antagonists.

                      Substance P is a naturally occurring undecapeptide  
belonging to the tachykinin family of peptides, the latter  
20    being named because of their prompt stimulatory action on  
smooth muscle tissue. More specifically, substance P is a  
pharmacologically active neuropeptide that is produced in  
mammals and possesses a characteristic amino acid sequence  
that is illustrated by D. F. Veber et al. in U.S. Patent No.  
25    4,680,283. The wide involvement of substance P and other  
tachykinins in the pathophysiology of numerous diseases has  
been amply demonstrated in the art. For instance, substance  
P has been shown to be involved in the transmission of pain  
or migraine (see B.E.B. Sandberg et al., Journal of  
30    Medicinal Chemistry, 25, 1009 (1982)), as well as in central  
nervous system disorders such as anxiety and schizophrenia,  
in respiratory and inflammatory diseases such as asthma and  
rheumatoid arthritis, respectively, in rheumatic diseases  
such as fibrositis, and in gastrointestinal disorders and  
35    diseases of the GI tract such as ulcerative colitis and  
Crohn's disease, etc. (see D. Regoli in "Trends in Cluster  
Headache," edited by F. Sicuteri et al., Elsevier Scientific  
Publishers, Amsterdam, pp. 85-95 (1987)).


                      Quinuclidine, piperidine, and azanorbornane derivatives  
40    and related compounds that exhibit activity as substance P  
receptor antagonists are referred to in United States Patent

-2-

Application 566,338 filed November 20, 1989, United States Patent Application 724,268, filed July 1, 1991, PCT Patent Application PCT/US 91/02853, filed April 25, 1991, PCT Patent Application PCT/US 91/03369, filed May 14, 1991, PCT Patent Application PCT/US 91/05776, filed August 20, 1991, PCT Patent Application PCT/US 92/00113, filed January 17, 1992, PCT Patent Application PCT/US 92/03571, filed May 5, 1992, PCT Patent Application PCT/US 92/03317, filed April 28, 1992, PCT Patent Application PCT/US 92/04697, filed June 11, 1992, United States Patent Application 766,488, filed September 26, 1991, United States Patent Application 790,934, filed November 12, 1991, PCT Patent Application PCT/US 92/04002, filed May 19, 1992, and Japanese Patent Application No. 065337/92, filed March 23, 1992.

# 15                    Summary of the Invention

The present invention relates to compounds of the formula



wherein ring A is an aryl group selected from phenyl, naphthyl, thienyl, dihydroquinolinyl and indolinyl, and wherein the side chain containing  $NR^2R^3$  is attached to a carbon atom of ring A;

W is hydrogen,  $(C_1-C_6)$ alkyl,  $S-(C_1-C_3)$ alkyl, halo or  $(C_1-C_6)$ alkoxy optionally substituted with from one to three fluorine atoms;

30                     $R^1$  is selected from amino,  $(C_1-C_6)$ alkylamino, di- $(C_1-C_6)$ alkylamino,  $-S(O)_v-(C_1-C_{10})$ -alkyl wherein v is zero, one or two,  $-S(O)_v$ -aryl wherein v is zero, one or two,  $-O$ -aryl,  $-SO_2NR^4R^5$  wherein each of  $R^4$  and  $R^5$  is, independently,  $(C_1-C_6)$ alkyl, or  $R^4$  and  $R^5$ , together with the nitrogen to which they are attached, form a saturated ring containing one nitrogen and from 3 to 6

-3-


carbons,  $\text{-NHC(=O)(C}_1\text{-C}_6\text{)alkyl}$ ,  $\text{-N[(C}_1\text{-C}_6\text{)alkyl]C(=O)(C}_1\text{-C}_6\text{)alkyl}$ ,

5  $\text{-NHCCF}_3$ ,  $\text{-N[(C}_1\text{-C}_6\text{)alkyl]CCF}_3$ ,  $\text{(C}_1\text{-C}_{10}\text{)alkyl-N-SO}_2\text{-(C}_1\text{-C}_{10}\text{)alkyl}$   
 wherein one or both of the alkyl moieties may optionally be  
 10 substituted with from one to three fluorine atoms,

$\text{-N(SO}_2\text{-(C}_1\text{-C}_{10}\text{)alkyl)}_2$  and  $\text{(C}_1\text{-C}_{10}\text{)alkyl-N-SO}_2\text{-aryl}$ ; and wherein  
 the aryl moieties of said  $\text{-S(O)}_2\text{-aryl}$ ,  $\text{-O-aryl}$  and

15  $\text{(C}_1\text{-C}_{10}\text{)alkyl-N-SO}_2\text{-aryl}$  are independently selected from  
 phenyl and benzyl and may optionally be substituted with  
 from one to three substituents independently selected from  
 $\text{(C}_1\text{-C}_4\text{)alkyl}$ ,  $\text{(C}_1\text{-C}_4\text{)alkoxy}$  and halo;

20 or  $\text{R}^1$  is a group having the formula



wherein  $a$  is 0, 1 or 2 and the asterisk represents a  
 position meta to the  $\text{R}^2\text{R}^3\text{NCH}_2$  side chain;

$\text{R}^2$  is hydrogen or  $\text{-CO}_2\text{(C}_1\text{-C}_{10}\text{)alkyl}$ ;


$\text{R}^3$  is selected from

30


35

40

5




II




III

10




IV




V

15




VI

20



VII


25



VIII

30

or



IX

35

-5-


wherein  $R^6$  and  $R^{10}$  are independently selected from furyl, thienyl, pyridyl, indolyl, biphenyl and phenyl, wherein said phenyl may optionally be substituted with one or two substituents independently selected from halo,  $(C_1-C_{10})$  alkyl optionally substituted with from one to three fluorine atoms,  $(C_1-C_{10})$  alkoxy optionally substituted with from one to three fluorine atoms, carboxy, benzyloxycarbonyl and  $(C_1-C_3)$  alkoxy-carbonyl;

$R^7$  is selected from  $(C_3-C_4)$  branched alkyl,  $(C_5-C_6)$  branched alkenyl,  $(C_5-C_7)$  cycloalkyl, and the radicals named in the definition of  $R^6$ ;

$R^8$  is hydrogen or  $(C_1-C_6)$  alkyl;

$R^9$  and  $R^{19}$  are independently selected from phenyl, biphenyl, naphthyl, pyridyl, benzhydryl, thienyl or furyl, and  $R^9$  and  $R^{19}$  may optionally be substituted with from one to three substituents independently selected from halo,  $(C_1-C_{10})$  alkyl optionally substituted with from one to three fluorine atoms and  $(C_1-C_{10})$  alkoxy optionally substituted with from one to three fluorine atoms;

$Y$  is  $(CH_2)_l$ , wherein  $l$  is an integer from one to three, or  $Y$  is a group of the formula



$Z$  is oxygen, sulfur, amino,  $(C_1-C_3)$ alkylamino or  $(CH_2)_n$ , wherein  $n$  is zero, one or two;

$x$  is zero, one or two;  
 $y$  is zero, one or two;  
 $z$  is three, four or five;  
 $o$  is two or three;  
 $p$  is zero or one;  
 $r$  is one, two or three;



-6-

the ring containing  $(CH_2)_2$  may contain from zero to three double bonds, and one of the carbon atoms of  $(CH_2)_2$  may optionally be replaced by oxygen, sulfur or nitrogen;

$R^{11}$  is thienyl, biphenyl or phenyl optionally substituted with one or two substituents independently selected from halo,  $(C_1-C_{10})$  alkyl optionally substituted with from one to three fluorine atoms and  $(C_1-C_{10})$  alkoxy optionally substituted with from one to three fluorine atoms;

10  $X$  is  $(CH_2)_q$ , wherein  $q$  is an integer from 1 to 6, and wherein any one of the carbon-carbon single bonds in said  $(CH_2)_q$  may optionally be replaced by a carbon-carbon double bond, and wherein any one of the carbon atoms of said  $(CH_2)_q$  may optionally be substituted with  $R^{14}$ , and wherein any one  
15 of the carbon atoms of said  $(CH_2)_q$  may optionally be substituted with  $R^{15}$ ;

$m$  is an integer from 0 to 8, and any one of the carbon-carbon single bonds of  $(CH_2)_m$ , wherein both carbon atoms of such bond are bonded to each other and to another  
20 carbon atom of the  $(CH_2)_m$  chain, may optionally be replaced by a carbon-carbon double bond or a carbon-carbon triple bond, and any one of the carbon atoms of said  $(CH_2)_m$  may optionally be substituted with  $R^{17}$ ;

$R^{12}$  is a radical selected from hydrogen,  $(C_1-C_6)$  straight  
25 or branched alkyl,  $(C_3-C_7)$  cycloalkyl wherein one of the carbon atoms may optionally be replaced by nitrogen, oxygen or sulfur; aryl selected from biphenyl, phenyl, indanyl and naphthyl; heteroaryl selected from thienyl, furyl, pyridyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, triazolyl, tetrazolyl and quinolyl; phenyl- $(C_2-C_6)$  alkyl, benzhydryl and  
30 benzyl, wherein the point of attachment on  $R^{12}$  is a carbon atom unless  $R^{12}$  is hydrogen, and wherein each of said aryl and heteroaryl groups and the phenyl moieties of said benzyl, phenyl- $(C_2-C_6)$  alkyl and benzhydryl may optionally be  
35 substituted with one or more substituents independently selected from halo, nitro,  $(C_1-C_{10})$  alkyl optionally

-7-

substituted with from one to three fluorine atoms, (C<sub>1</sub>-C<sub>10</sub>)  
alkoxy optionally substituted with from one to three  
fluorine atoms, amino, hydroxy-(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
(C<sub>1</sub>-C<sub>6</sub>)alkoxy-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)-alkylamino,

5 (C<sub>1</sub>-C<sub>6</sub>)alkyl-O-C(=O)-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-O-C(=O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl,

10 (C<sub>1</sub>-C<sub>6</sub>)alkyl-C(=O)-O-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-C(=O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl-O-,

15 (C<sub>1</sub>-C<sub>6</sub>)alkyl-C(=O)-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-C(=O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl-,

20 di-(C<sub>1</sub>-C<sub>6</sub>)alkylamino, -C(=O)NH-(C<sub>1</sub>-C<sub>6</sub>)alkyl,

(C<sub>1</sub>-C<sub>6</sub>)-alkyl-C(=O)NH-(C<sub>1</sub>-C<sub>6</sub>)alkyl, -NHCH(=O) and -NHC(=O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl;  
and wherein one of the phenyl moieties of said benzhydryl  
25 may optionally be replaced by naphthyl, thienyl, furyl or  
pyridyl;

R<sup>13</sup> is hydrogen, phenyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl;


or R<sup>12</sup> and R<sup>13</sup>, together with the carbon to which they  
are attached, form a saturated carbocyclic ring having from  
30 3 to 7 carbon atoms wherein one of said carbon atoms that is  
neither the point of attachment of the spiro ring nor  
adjacent to such point of attachment may optionally be  
replaced by oxygen, nitrogen or sulfur;

R<sup>14</sup> and R<sup>15</sup> are each independently selected from  
35 hydrogen, hydroxy, halo, amino, oxo (=O), cyano, hydroxy-(C<sub>1</sub>-  
C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino,

40 di-(C<sub>1</sub>-C<sub>6</sub>)alkylamino, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, -C(=O)OH,

45 (C<sub>1</sub>-C<sub>6</sub>)alkyl-O-C(=O)-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-O-C(=O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl,

-8-



5  $(C_1-C_6)\text{alkyl}-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-$ ,  $(C_1-C_6)\text{alkyl}-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-(C_1-C_6)\text{alkyl}-$ , and the radicals set forth in the definition of  $R^{12}$ ;

10

$R^{16}$  is  $\text{NH}\overset{\overset{\text{O}}{\parallel}}{\text{C}}\text{R}^{18}$ ,  $\text{NHCH}_2\text{R}^{18}$ ,  $\text{SO}_2\text{R}^{18}$ ,  $\text{CO}_2\text{H}$  or one of the radicals set forth in any of the definitions of  $R^{12}$ ,  $R^{14}$  and  $R^{15}$ ;

15

$R^{17}$  is oximino ( $=\text{NOH}$ ) or one of the radicals set forth in any of the definitions of  $R^{12}$ ,  $R^{14}$  and  $R^{15}$ ; and

$R^{18}$  is  $(C_1-C_6)\text{alkyl}$ , hydrogen, phenyl or phenyl  $(C_1-C_6)\text{alkyl}$ ;

20

with the proviso that (a) when  $m$  is 0, one of  $R^{16}$  and  $R^{17}$  is absent and the other is hydrogen, (b) when  $R^3$  is a group of the formula VIII,  $R^{14}$  and  $R^{15}$  cannot be attached to the same carbon atom, (c) when  $R^{14}$  and  $R^{15}$  are attached to the same carbon atom, then either each of  $R^{14}$  and  $R^{15}$  is independently selected from hydrogen, fluoro,  $(C_1-C_6)\text{alkyl}$ , hydroxy- $(C_1-C_6)\text{alkyl}$  and  $(C_1-C_6)\text{alkoxy}-(C_1-C_6)\text{alkyl}$ , or  $R^{14}$  and  $R^{15}$ , together with the carbon to which they are attached, form a  $(C_3-C_6)$  saturated carbocyclic ring that forms a spiro compound with the nitrogen-containing ring to which they are attached; (d) when  $R^1$  is amino,  $(C_1-C_6)\text{alkylamino}$ , di- $(C_1-$

30

$\overset{\overset{\text{O}}{\parallel}}{\text{C}}_6)\text{alkylamino}$  or  $\text{NHC}(\overset{\overset{\text{O}}{\parallel}}{\text{C}}_1-C_6)\text{alkyl}$ ,  $R^3$  is a group of the formula II, III, IV, V or VI, and (e) when  $R^{14}$  or  $R^{15}$  is attached to a carbon atom of X or  $(\text{CH}_2)_y$ , that is adjacent to the ring nitrogen, then  $R^{14}$  or  $R^{15}$ , respectively, must be a substituent wherein the point of attachment is a carbon atom.

35

The present invention also relates to the pharmaceutically acceptable acid addition and base salts of compounds of the formula I. The acids which are used to prepare the pharmaceutically acceptable acid addition salts

40

-9-

of the aforementioned base compounds of this invention are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, 5 bisulfate, phosphate, acid phosphate, acetate, lactate, citrate, acid citrate, tartrate, bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and pamoate [i.e., 1,1'-methylene-10 bis-(2-hydroxy-3-naphthoate)]salts. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this invention are those which form non-toxic base salts with the acidic compounds of formula I. Such non-toxic base salts include those derived from such 15 pharmacologically acceptable cations as sodium, potassium calcium and magnesium, etc.

The term "halo", as used herein, unless otherwise indicated, includes chloro, fluoro, bromo and iodo.

The term "alkyl", as used herein, unless otherwise 20 indicated, includes saturated monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof.

The term "alkoxy", as used herein, includes O-alkyl groups wherein "alkyl" is defined as above.

25 The term "one or more substituents," as used herein, includes from one to the maximum number of substituents possible based on the number of available bonding sites.

Preferred compounds of the formula I include those wherein the substituents at positions "2" and "3" of the 30 nitrogen containing ring of R<sup>3</sup> are in a cis configuration. When R<sup>3</sup> is a group of the formula VII or VIII, "a cis configuration," as used herein, means that the non-hydrogen substituent at position "3" is cis to R<sup>12</sup>.

Other preferred compounds of the formula I are those 35 wherein R<sup>3</sup> is a group of the formula II, III, VII or IX; R<sup>2</sup> is hydrogen; ring A is phenyl or indoliny1; W is (C<sub>1</sub>-C<sub>3</sub>)alkoxy optionally substituted with from one to five

-10-

fluorine atoms; and  $R^1$  is  $S(O)_v-(C_1-C_{10})$ alkyl wherein  $v$  is zero, one or two,  $S(O)_v$ -aryl wherein  $v$  is zero,

5 one or two,  $-O$ -aryl,  $(C_1-C_{10})$ alkyl- $N-SO_2-(C_1-C_{10})$ alkyl wherein one or both of the alkyl moieties may optionally be substituted with from one to three fluorine atoms,  $-N(SO_2-$

10  $(C_1-C_{10})$ alkyl) $_2$  or  $(C_1-C_{10})$ alkyl- $N-SO_2$ -aryl wherein said aryl is phenyl or benzyl and may optionally be substituted with from one to three substituents independently selected from  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$ alkoxy and halo.

More preferred compounds of the formula I are the  
15 foregoing preferred compounds wherein: (a)  $R^3$  is a group of the formula III and  $R^9$  is benzhydryl; (b)  $R^3$  is a group of the formula VII, each of  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$  and  $R^{16}$  is hydrogen,  $m$  is zero and  $X$  is  $-(CH_2)_3-$ ; or (c)  $R^3$  is a group of the formula IX,  $r$  is two and  $R^{19}$  is benzhydryl.

20 Other more preferred compounds of the formula I are those wherein: (a)  $R^3$  is a group of the formula III wherein the substituents at positions "2" and "3" of the nitrogen containing ring are in the cis configuration,  $R^9$  is benzhydryl and ring A is phenyl; (b)  $R^3$  is a group of the  
25 formula VII wherein  $R^{12}$  and the substituent at position "3" of the nitrogen containing ring are in the cis configuration, ring A is phenyl,  $R^{12}$  is phenyl, each of  $R^2$ ,  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$  and  $R^{16}$  is hydrogen,  $m$  is zero,  $W$  is methoxy or isopropoxy,  $X$  is  $-(CH_2)_3-$  and  $R^1$  is  $S(O)_v-(C_1-C_{10})$ alkyl wherein  
30

$v$  is zero, one or two, or  $(C_1-C_{10})$ alkyl- $N-SO_2-(C_1-C_{10})$ alkyl; or  
(c)  $R^3$  is a group of the formula IX wherein the substituents at positions "2" and "3" of the nitrogen containing ring are  
35 in the cis configuration,  $R^{19}$  is benzhydryl,  $r$  is two and ring A is phenyl.

Especially preferred compounds of the formula I are those wherein  $R^3$  is a group of the formula III,  $R^9$  is benzhydryl, ring A is phenyl,  $W$  is selected from  $OCF_3$ ,

-11-

isopropoxy,  $\text{OCH}_3$ ,  $\text{OCHF}_2$  and  $\text{OCH}_2\text{CF}_3$ , and  $\text{R}^1$  is selected from amino,  $(\text{C}_1\text{-C}_6)\text{alkylamino}$ , di- $(\text{C}_1\text{-C}_6)\text{alkylamino}$ , and  $-\text{S}(\text{O})_v-(\text{C}_1\text{-C}_{10})\text{alkyl}$  wherein  $v$  is zero, one or two.

Other especially preferred compounds of this invention are those wherein  $\text{R}^3$  is a group of the formula VII, each of  $\text{R}^{13}$ ,  $\text{R}^{14}$ ,  $\text{R}^{15}$  and  $\text{R}^{16}$  is hydrogen,  $m$  is zero,  $X$  is  $-(\text{CH}_2)_3-$ , ring A is phenyl, W is selected from  $\text{OCF}_3$ ,  $\text{OCH}_3$ , isopropoxy,  $\text{OCHF}_2$  and  $\text{OCH}_2\text{CF}_3$ , and  $\text{R}^1$  is selected from  $-\text{S}(\text{O})_v-(\text{C}_1\text{-C}_{10})\text{alkyl}$

wherein  $v$  is zero, one or two, and  $(\text{C}_1\text{-C}_{10})\text{alkyl-N-SO}_2-(\text{C}_1\text{-C}_{10})\text{alkyl}$ .

Specific preferred compounds of the formula I include the following:

- 15 (2S,3S)-3-[2-methoxy-5-(N-acetyl-N-methylamino)benzylamino]-2-diphenylmethyl-1-azabicyclo[2.2.2]octane; -  
(1SR, 2SR, 3SR, 4RS)-3-(2-methoxy-5-(N-methyl-N-trifluoromethane-sulfonylamino)benzyl)amino-2-benzhydryl-[2.2.1]-azanorbornane;
- 20 (1SR,2SR,3SR,4RS)-3-[2-methoxy-5-(N-thiazolidine-S,S-dioxide)benzyl]amino-2-benzhydryl-[2.2.1]-1-azanorbornane;  
(1SR,2SR,3SR,4RS)-3-[(2,3-dihydro-5-methoxy-1-methanesulfonyl-6-indolyl)methylamino]-2-benzhydryl-[2.2.1]-1-azanorbornane;
- 25 (2S,3S)-3-(2-methoxy-5-methylthiobenzyl)amino-2-phenylpiperidine;  
(2S,3S)-3-(2-methoxy-5-methylsulfonylbenzyl)amino-2-phenylpiperidine;  
(2S,3S)-3-[2-methoxy-5-(N-methyl-N-methanesulfonylamino)-benzyl]amino-2-phenylpiperidine;
- 30 (2S,3S)-3-[2-trifluoromethoxy-5-(N-methyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine;  
(2S,3S)-3-[2-isopropoxy-5-(N-methyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- 35 (2S,3S)-3-[2-methoxy-5-(N-isopropyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine;  
(2S,3S)-3-[2-isopropoxy-5-(N-isopropyl-N-methanesulfonyl-amino)benzyl]amino-2-phenylpiperidine;

-12-

- (2S,3S)-3-[2-isopropoxy-5-(N-isopropyl-N-methanesulfonyl-amino)benzyl]amino-2-phenylpiperidine;
- (2S,3S)-3-[2-methoxy-5-(N-cyclopentyl-N-methanesulfonyl-amino)benzyl]amino-2-phenylpiperidine;
- 5 (2S,3S)-3-[2-methoxy-5-(N-methyl-N-trifluoromethanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- (2S,3S)-3-[2-isopropoxy-5-(N-methyl-N-trifluoromethanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- (2S,3S)-3-[2-methoxy-5-(N-methyl-N-isopropylsulfonylamino)-benzyl]amino-2-phenylpiperidine;
- 10 (2S,3S)-3-[2-methoxy-5-(N-thiazolidine-S,S-dioxide)-benzyl]amino-2-phenyl-piperidine;
- (2S,3S)-3-[(2,3-dihydro-5-methoxy-1-methanesulfonyl-6-indolyl)methylamino]-2-phenylpiperidine;
- 15 (2S,3S)-3-[(2,3-dihydro-5-methoxy-2-methyl-1-methanesulfonyl-6-indolyl)methylamino]-2-phenylpiperidine;
- (2SR,3SR,4RS)-2-benzhydryl-4-(2-hydroxyethyl)-3-(2-methoxy-5-methylthiobenzyl)aminopyrrolidine;
- (2SR,3SR,4RS)-2-benzhydryl-4-(2-hydroxyethyl)-3-(2-methoxy-5-(N-methyl-N-methanesulfonylamino)benzyl)amino-
- 20 pyrrolidine;
- (2SR,3SR,4RS)-2-benzhydryl-4-(2-hydroxyethyl)-3-(2-methoxy-5-(N,thiazolidine-S,S-dioxide)benzyl)aminopyrrolidine;
- 25 (2S,3S)-N-(2-methoxy-5-methylthiophenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(2-methoxy-5-dimethylaminophenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(5-ethylthio-2-methoxyphenyl)methyl-2-
- 30 diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(5-trifluoroacetyl-amino-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(5-amino-2-methoxyphenyl)methyl-2-diphenyl-
- 35 methyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(2-methoxy-5-methylsulfinylphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine; and

-13-

(2S,3S)-N-(2-methoxy-5-methylsulfonylphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine.

Other compounds of the formula I include:

- 4-(2-methylthiophenyl)methylamino-3-phenyl-2-  
5 azabicyclo[3.3.0]octane;  
4-benzhydryl-5-(2-methylthiophenyl)methylamino-3-  
azabicyclo[4.1.0]heptane;  
4-(2-methylthiophenyl)methylamino-3-phenyl-2-  
azabicyclo[4.4.0]decane;  
10 8-benzhydryl-7-(2-methylthio-5-trifluoromethoxy-  
phenyl)methylamino-9-azatricyclo[4.3.1.0<sup>4,9</sup>]decane;  
9-benzhydryl-8-(2-methylthio-5-trifluoromethoxy-  
phenyl)methylamino-10-azatricyclo[4.4.1.0<sup>5,10</sup>]undecane;  
9-benzhydryl-8-(2-methylthio-5-trifluoromethoxy-  
15 phenyl)methylamino-3-thia-10-azatricyclo[4.4.1.0<sup>5,10</sup>]undecane;  
2-benzhydryl-3-(2-methylthiophenyl)methylamino-5,6-  
pentamethylene-quinuclidine;  
2-benzhydryl-3-(2-methylthiophenyl)methylamino-5,6-  
trimethylene-quinuclidine;  
20 cis-3-(2-phenoxyphenyl)methylamino-2-benzhydryl-  
quinuclidine;  
8-benzhydryl-9-(2-methylthiophenyl)methylamino-7-  
azatricyclo[4.4.1.0<sup>5,10</sup>]undecane;  
2-benzhydryl-3-(2-methylthiophenyl)methylamino-1-  
25 azabicyclo[3.2.2]nonane;  
2-benzhydryl-3-(2-methylthiophenyl)methylamino-1-  
azabicyclo[2.2.1]heptane;  
3-(2-methylthiophenyl)methylamino-2-phenyl-1-  
azabicyclo[2.2.1]heptane;  
30 N-[3-(4-benzhydryl-1-azabicyclo[2.2.1]hept-3-  
ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-  
isopropylsulfonamide;  
N-[3-(7-benzhydryl-1-azabicyclo[3.2.2]non-6-  
ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonamide;  
35 N-[3-(7-benzhydryl-1-azabicyclo[3.2.2]non-6-  
ylaminomethyl)-4-trifluoromethoxyphenyl]-N-methyl-  
methanesulfonamide;



-14-

N-[3-(8-benzhydryl-1-azabicyclo[4.2.2]dec-7-ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonamide;

N-[3-(8-benzhydryl-1-azabicyclo[4.2.2]dec-7-ylaminomethyl)-4-trifluoromethoxyphenyl]-N-methyl-  
5 methanesulfonamide;

N-[3-(4-benzhydryl-1-azabicyclo[2.2.1]hept-3-ylaminomethyl)-4-methoxyphenyl]-N-methyl-isopropylsulfonamide;

N-[3-(4-benzhydryl-1-azabicyclo[2.2.1]hept-3-ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-  
10 trifluoromethanesulfonamide;

N-[3-(4-benzhydryl-1-azabicyclo[2.2.1]hept-3-ylaminomethyl)-4-methoxyphenyl]-isopropylsulfone;

N-[3-(4-benzhydryl-1-azabicyclo[2.2.1]hept-3-ylaminomethyl)-4-methoxyphenyl]-methylsulfone;  
15

N-[3-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonamide;

N-[3-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-ylaminomethyl)-4-isopropoxyphenyl]-N-methyl-  
20 methanesulfonamide;

N-[3-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-ylaminomethyl)-4-trifluoromethoxyphenyl]-N-methyl-methanesulfonamide;

N-[3-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-methanesulfonamide;  
25

N-[3-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-ylaminomethyl)-4-methoxyphenyl]-N-methyl-trifluoromethanesulfonamide;

N-[3-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-ylaminomethyl)-4-methoxyphenyl]-methanesulfone;  
30

N-[3-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-ylaminomethyl)-4-trifluoromethoxyphenyl]-methanesulfone;

N-[3-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-ylaminomethyl)-4-methoxyphenyl]-isopropylsulfone;  
35

N-[3-(9-benzhydryl-1-azatricyclo[5.2.2.0<sup>2,6</sup>]undec-8-ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonamide;

-15-

N-[3-(9-benzhydryl-1-azatricyclo[5.2.2.0<sup>2,6</sup>]undec-8-ylaminomethyl)-4-isopropoxyphenyl]-N-methyl-methanesulfonamide;

5 N-[3-(9-benzhydryl-1-azatricyclo[5.2.2.0<sup>2,6</sup>]undec-8-ylaminomethyl)-4-trifluoromethoxyphenyl]-N-methyl-methanesulfonamide;

N-[3-(10-benzhydryl-octahydro-1,4-ethano-quinolin-9-ylaminomethyl)-4-isopropoxyphenyl]-N-isopropyl-methanesulfonamide;

10 N-[3-(9-benzhydryl-1-azatricyclo[5.2.2.0<sup>2,6</sup>]undec-8-ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-methanesulfonamide;

15 N-[3-(9-benzhydryl-1-azatricyclo[5.2.2.0<sup>2,6</sup>]undec-8-ylaminomethyl)-4-methoxyphenyl]-N-methyl-isopropyl-sulfonamide;

N-[3-(9-benzhydryl-1-azatricyclo[5.2.2.0<sup>2,6</sup>]undec-8-ylaminomethyl)-4-methoxyphenyl]-methanesulfone;

N-[3-(9-benzhydryl-1-azatricyclo[5.2.2.0<sup>2,6</sup>]undec-8-ylaminomethyl)-4-trifluoromethoxyphenyl]-methanesulfone;

20 N-[3-(9-benzhydryl-1-azatricyclo[5.2.2.0<sup>2,6</sup>]undec-8-ylaminomethyl)-4-methoxyphenyl]-isopropylsulfone;

N-[3-(9-benzhydryl-8-azatricyclo[5.3.1.0<sup>3,8</sup>]undec-10-ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonamide;

25 N-[3-(10-benzhydryl-9-azatricyclo[6.3.1.0<sup>3,9</sup>]dodec-11-ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonamide;

N-[3-(9-benzhydryl-8-azatricyclo[5.3.1.0<sup>3,8</sup>]undec-10-ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-methanesulfonamide;

30 N-[3-(10-benzhydryl-9-azatricyclo[6.3.1.0<sup>3,9</sup>]dodec-11-ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-methanesulfonamide;

N-[3-(10-benzhydryl-9-azatricyclo[6.3.1.0<sup>3,9</sup>]dodec-11-ylaminomethyl)-4-methoxyphenyl]-N-methyl-isopropyl-sulfonamide;

35 N-[3-(11-benzhydryl-1-azatricyclo[6.3.1.0<sup>3,9</sup>]dodec-10-ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonamide;

-16-

N-[3-(11-benzhydryl-1-azatricyclo[6.3.1.0<sup>3,9</sup>]dodec-10-ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-methanesulfonamide;

5 N-[3-(11-benzhydryl-1-azatricyclo[6.3.1.0<sup>3,9</sup>]dodec-10-ylaminomethyl)-4-methoxyphenyl]-N-methyl-isopropyl-sulfonamide;

N-[3-(11-benzhydryl-1-azatricyclo[6.3.1.0<sup>3,9</sup>]dodec-10-ylaminomethyl)-4-methoxyphenyl]-N-methyl-trifluoromethane-sulfonamide;

10 N-[3-(3-benzhydryl-octahydro-2,5-methano-isoquinolin-4-ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonamide;

N-[3-(3-benzhydryl-octahydro-2,5-methano-isoquinolin-4-ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-methane-sulfonamide;

15 (2-trifluoromethoxy-5-methylsulfonylbenzyl)-(2-phenylpiperidin-3-yl)-amine;

(2-difluoromethoxy-5-methylsulfonylbenzyl)-(2-phenylpiperidin-3-yl)-amine;

20 (2-cyclopropoxy-5-methylsulfonylbenzyl)-(2-phenylpiperidin-3-yl)-amine;

(2-cyclopentyloxy-5-methylsulfonylbenzyl)-(2-phenylpiperidin-3-yl)-amine;

(2-isopropoxy-5-methylsulfonylbenzyl)-(2-phenylpiperidin-3-yl)-amine;

25 (2-methoxy-5-isopropylsulfonylbenzyl)-(2-phenylpiperidin-3-yl)-amine;

N-[4-methoxy-3-(3-phenyl-decahydro-isoquinolin-4-ylaminomethyl)-phenyl]-N-methyl-methanesulfonamide;

30 N-[4-trifluoromethoxy-3-(3-phenyl-decahydroisoquinolin-4-ylaminomethyl)-phenyl]-N-methyl-methanesulfonamide;

N-[4-methoxy-3-(3-phenyl-decahydro-isoquinolin-4-ylaminomethyl)-phenyl]-N-isopropyl-methanesulfonamide;

N-[4-methoxy-3-(2-phenyl-decahydro-quinolin-3-ylaminomethyl)-phenyl]-N-methyl-methanesulfonamide;

35 N-[4-methoxy-3-(2-phenyl-decahydro-quinolin-3-ylaminomethyl)-phenyl]-N-isopropyl-methanesulfonamide;

-17-


- N-[4-methoxy-3-(2-phenyl-decahydro-quinolin-3-ylaminomethyl)-phenyl]-N-methyl-isopropylsulfonamide;
- N-[4-methoxy-3-(2-phenyl-octahydro-[1]pyrindin-3-ylaminomethyl)-phenyl]-N-methyl-methanesulfonamide;
- 5 N-[4-methoxy-3-(2-phenyl-octahydro-[1]pyrindin-3-ylaminomethyl)-phenyl]-N-isopropyl-methanesulfonamide;
- N-[4-methoxy-3-(2-phenyl-octahydro-[1]pyrindin-3-ylaminomethyl)-phenyl]-N-methyl-trifluoromethanesulfonamide;
- 10 [b]pyridin-3-ylaminomethyl)-phenyl]-N-methyl-methanesulfonate;
- N-[4-methoxy-3-(2-phenyl-octahydro-indol-3-ylaminomethyl)-phenyl]-N-methyl-methanesulfonate;
- N-[3-(2-benzhydryl-decahydro-cyclohepta[b]pyridin-3-ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonate;
- 15 N-[3-(7-benzhydryl-1-aza-bicyclo[3.2.1]oct-6-ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonamide;
- N-[3-(7-benzhydryl-1-aza-bicyclo[3.2.1]oct-6-ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-
- 20 methanesulfonamide;
- N-[3-(7-benzhydryl-1-aza-bicyclo[3.2.1]oct-6-ylaminomethyl)-4-methoxyphenyl]-N-methyl-isopropyl-sulfonamide;
- N-[3-(7-benzhydryl-1-aza-bicyclo[3.2.1]oct-6-ylaminomethyl)-4-methoxyphenyl]-N-methyl-trifluoromethane-
- 25 sulfonamide;
- N-[3-(8-benzhydryl-1-aza-bicyclo[4.2.1]non-7-ylaminomethyl)-4-methoxyphenyl]-N-methyl-methanesulfonamide;
- N-[3-(8-benzhydryl-1-aza-bicyclo[4.2.1]non-7-ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-
- 30 methanesulfonamide;
- N-[3-(9-benzhydryl-1-aza-bicyclo[5.2.1]dec-8-ylaminomethyl)-4-methoxyphenyl]-N-isopropyl-methanesulfonamide;
- 35 N-[3-(9-benzhydryl-1-aza-bicyclo[5.2.1]dec-8-ylaminomethyl)-4-methoxyphenyl]-N-methyl-isopropyl-sulfonamide;

-18-

N-[3-(9-benzhydryl-1-aza-bicyclo[5.2.1]dec-8-ylaminomethyl)-4-methoxyphenyl]-methanesulfone; and


N-[3-(9-benzhydryl-1-aza-bicyclo[5.2.1]dec-8-ylaminomethyl)-4-trifluoromethoxyphenyl]-methanesulfone.

5 The present invention also relates to compounds of the formulae



XI

and



XII

wherein ring A, R<sup>1</sup>, R<sup>3</sup> and W are defined as above. These  
25 compounds are intermediate in the synthesis of compounds of the formula I.

The present invention also relates to a pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g.,  
30 arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic  
35 diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such

-19-

as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic  
5 neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable  
10 salt thereof, effective in treating or preventing such condition, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis,  
15 psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and  
20 Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia,  
25 neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including  
30 a human, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such condition.

The present invention also relates to a pharmaceutical  
35 composition for antagonizing the effects of substance P in a mammal, including a human, comprising a substance P antagonizing amount of a compound of the formula I, or a

-20-

pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of antagonizing the effects of substance P in a mammal, including a human, comprising administering to said mammal a substance P antagonizing amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder in a mammal, including a human, resulting from an excess of substance P, comprising a substance P antagonizing amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a disorder in a mammal, including a human, resulting from an excess of substance P, comprising administering to said mammal a substance P antagonizing amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof.

The present invention also relates to a pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fasciitis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus

-21-

erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of  
5 substance P at its receptor site, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis,  
10 psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and  
15 Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia,  
20 neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and  
25 a human, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance P at its receptor site.

The present invention also relates to a pharmaceutical  
30 composition for treating or preventing a disorder in a mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable  
35 salt thereof, effective in antagonizing the effect of substance P at its receptor site, and a pharmaceutically acceptable carrier.



-22-

The present invention also relates to a method of treating or preventing a disorder in mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance P at its receptor site.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder in a mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such disorder, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a disorder in mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such disorder.


The compounds of the formula I have chiral centers and therefore exist in different enantiomeric forms. This invention relates to all optical isomers and all stereoisomers of compounds of the formula I, and mixtures thereof.

-23-


Detailed Description of the Invention

The compounds of the formula I may be prepared as described in the following reaction schemes and discussion. Unless otherwise indicated, ring A, W, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>,  
5 R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup>, R<sup>19</sup>, X, Z, Y, m, n, o, p, q, r, x, y, and z, and structural formulas I, II, III, IV, V, VI, VII, VIII, IX, XI and XII in the reaction schemes and discussion that follow are defined as above.



-24-

Scheme 1

-25-

Scheme 2

-26-


Scheme 3Scheme 4

-27-

Scheme 1 illustrates the preparation of compounds of the formula I from starting materials of the formula X wherein G is hydrogen, hydroxy, chloro, bromo or (C<sub>1</sub>-C<sub>6</sub>)alkoxy.

5 Referring to scheme 1, a compound of the formula X wherein G is hydrogen may be converted directly into the corresponding compound of the formula I by reacting it with a compound of the formula NH<sub>2</sub>R<sup>3</sup> in the presence of a reducing agent. Reducing agents that may be used include sodium  
10 cyanoborohydride, sodium triacetoxyborohydride, sodium borohydride, hydrogen and a metal catalyst, zinc and hydrochloric acid, and formic acid. This reaction is typically conducted in a reaction inert solvent at a temperature from about 0°C to about 150°C. Suitable  
15 reaction inert solvents include lower alcohols (e.g., methanol, ethanol and isopropanol), 1,2-dichloroethane, acetic acid and tetrahydrofuran (THF). Preferably, the solvent is acetic acid, the temperature is about 25°C, the reducing agent is sodium triacetoxyborohydride, and the  
20 reaction is conducted in the presence of a dehydrating agent such as molecular sieves.

Alternatively, the reaction of a compound of the formula X with a compound of the formula NH<sub>2</sub>R<sup>3</sup> may be carried out in the presence of a dehydrating agent or using an  
25 apparatus designed to remove azeotropically the water generated, to produce an imine of the formula



XI

35 which is then reacted with a reducing agent as described above, preferably with sodium triacetoxyborohydride in an acetic acid or 1,2-dichloroethane solvent at about room

-28-

temperature. The preparation of the imine is generally carried out in a reaction inert solvent such as benzene, xylene or toluene, preferably toluene, at a temperature from about 25°C to about 110°C, preferably at about the reflux temperature of the solvent. Suitable dehydrating agents/solvent systems include titanium tetrachloride/dichloromethane titanium isopropoxide/dichloromethane and molecular sieves/THF. Titanium tetrachloride/dichloromethane is preferred.

10 Compounds of the formula X wherein G is hydroxy, chloro, bromo or (C<sub>1</sub>-C<sub>6</sub>)alkoxy may be converted into the corresponding compounds of formula XII having the desired R<sup>3</sup> group by reacting them with the appropriate compound of the formula NH<sub>2</sub>R<sup>3</sup> under conditions that will be obvious to those skilled in the art, and then reducing the resulting amides to yield the desired compounds having formula I wherein R<sup>2</sup> is hydrogen. When G is hydroxy, the compound of formula X is reacted with NH<sub>2</sub>R<sup>3</sup> in the presence of an activating agent. Appropriate activating agents include carbonyldiimidazole, diethylphosphoryl cyanide and dicyclohexylcarbodiimide. Carbonyldiimidazole is preferred. This reaction is generally conducted at a temperature from about 0°C to about 50°C, preferably at about 25°C, in an inert solvent such as chloroform, diethyl ether, THF or dimethylformamide (DMF).

25 When G is chloro or bromo, the reaction of the compound of formula X with the appropriate compound of formula NH<sub>2</sub>R<sup>3</sup> is typically carried out in the presence of an acid scavenger in an aprotic solvent at a temperature from about 0°C to about 100°C. Suitable acid scavengers include triethylamine (TEA), pyridine and inorganic salts such as sodium and potassium carbonate. Suitable solvents include methylene chloride (CH<sub>2</sub>Cl<sub>2</sub>), chloroform (CHCl<sub>3</sub>), benzene, toluene and tetrahydrofuran (THF). Preferably, the reaction is conducted in CH<sub>2</sub>Cl<sub>2</sub> at room temperature using TEA as the acid scavenger.

When G is O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, the reaction of the compound of formula NH<sub>2</sub>R<sup>3</sup> is usually conducted in an aprotic solvent

-29-

such as benzene, toluene, chlorobenzene or xylenes, at a temperature from about 25°C to about 100°C, preferably at about the reflux temperature of the solvent.

Reduction of the compound of formula XII so formed yields the corresponding compound of the formula I wherein  $R^2$  is hydrogen. This is generally accomplished using a reducing agent such as lithium aluminum hydride, borane dimethylsulfide complex or diborane, in an aprotic solvent such as THF, dioxane or diethyl ether, at a temperature from about 0°C to about 70°C. Preferably, the reducing agent is borane dimethylsulfide complex and the reaction is carried out at about room temperature in an ethereal solvent such as THF.

Compounds of the formula I wherein  $R^2$  is hydrogen may be converted into the corresponding compounds wherein  $R^2$  is  $-CO_2(C_1-C_{10})$ alkyl by reacting them with a  $(C_1-C_{10})$ alkyl halo carbonate such as methyl or ethyl chloroformate in the presence of an acid scavenger. Typically, this reaction is conducted in an polar solvent such as chloroform, methylene chloride, water or a water/acetone mixture, at a temperature from about 0°C to about 100°C, preferably at about room temperature. Suitable acid scavengers include triethylamine, pyridine and potassium and sodium carbonate or bicarbonate.

When  $R^3$  is a group of the formula II, the starting materials of the formula  $NH_2R^3$  may be prepared as described in United States Patent Application Serial No. 566,338, filed July 20, 1990. This application is incorporated herein in its entirety.

When  $R^3$  is a group of the formula III, the starting materials of the formula  $NH_2R^3$  may be prepared as described in United States Patent Application Serial No. 532,525, filed June 1, 1990 and PCT Patent Application PCT/US 91/02853, filed April 25, 1991. Both these applications are incorporated herein in their entirety.

When  $R^3$  is a group of the formula IV, V or VI, the starting materials of the formula  $NH_2R^3$  may be prepared as



When R<sup>3</sup> is a group of the formula VIII, the starting materials of the formula NH<sub>2</sub>R<sup>3</sup> may be prepared as described in PCT Patent Application PCT/US 91/05776, filed August 20, 1991, United States Patent Application Serial No. 800,667, filed November 27, 1991 and PCT Patent Application PCT/US 92/00065, filed January 14, 1992. These applications are incorporated herein in their entirety.

Scheme 2 illustrates the preparation of the starting materials of formula X wherein G is hydrogen and R<sup>1</sup> is

Referring to scheme 2, a compound of the formula XIII

35 wherein R<sup>1</sup> is other than -NHCCF<sub>3</sub>, or -SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup> is reacted with titanium tetrachloride (TiCl<sub>4</sub>) and dichloromethyl methyl ether (CHCl<sub>2</sub>-O-CH<sub>3</sub>) at a temperature from about 0°C to about room temperature in a methylene chloride solvent to yield

-31-

the corresponding aldehyde of formula X wherein G is hydrogen. Alternatively, the compound of the formula XIII may be reacted with hexamethylene tetraamine and trifluoroacetic acid at about 70°C to yield the same  
5 product.

Those compounds of the formula XIII wherein R<sup>1</sup> is -SO-(C<sub>1</sub>-C<sub>10</sub>)alkyl, -SO<sub>2</sub>-(C<sub>1</sub>-C<sub>10</sub>)alkyl, -SO-aryl or -SO<sub>2</sub>-aryl may be obtained from their deoxygenated counterparts of the formula XIII wherein R<sup>1</sup> is -S-(C<sub>1</sub>-C<sub>10</sub>)alkyl or -S-aryl by reacting  
10 them with an oxidizing agent. For example, such oxidation may be carried out using metachloroperbenzoic acid in methylene chloride at about room temperature. It may also be carried out using peroxyphthalic acid magnesium hydrate in aqueous ethanol at a temperature from about 70°C to about  
15 100°C. The foregoing oxidation reactions produce mixtures of the oxy and dioxy products (-SO- and -SO<sub>2</sub>-) which can be separated by ordinary means.

Compounds of the formula X wherein G is hydrogen and R<sup>1</sup> is -NHCOCF<sub>3</sub> may be obtained using procedures known to those  
20 skilled in the art. Scheme 3 illustrates one method of preparing such compounds. Referring to scheme 3, the -CHO group of a nitro benzaldehyde of the formula XIV is protected by conversion to the corresponding 1,3-dioxolane of formula XV. This reaction is generally carried out by  
25 heating a mixture of the nitrobenzaldehyde and ethylene glycol in an inert solvent such as benzene or toluene, preferably in the presence of an acid such as p-toluenesulfonic acid, and preferably at the reflux temperature of the solvent to remove the water formed in the  
30 reaction. The resulting compound of formula XV is then treated with hydrogen gas and a metal catalyst such as palladium on carbon in a reaction inert solvent such as ethyl acetate or a lower alcohol to convert the NO<sub>2</sub> group to an NH<sub>2</sub> group and produce the corresponding compound of  
35 formula XVI.

The resulting intermediate of formula XVI is then acylated with a reagent such as ethyl trifluoroacetate in

-32-

methanol or trifluoroacetic anhydride in methylene chloride at a temperature from about 0°C to about 50°C, preferably at about room temperature, to produce the corresponding trifluoroacetamide of the formula XVII. Treatment of this  
5 amide with a mixture of aqueous hydrochloric acid in acetone at a temperature from about 0°C to about 50°C, preferably at room temperature, will convert the dioxolane to the desired compound of formula X wherein R<sup>1</sup> is NHCOCF<sub>3</sub> and G is hydrogen.

10 Scheme 4 illustrates the preparation of the starting materials of the formula X wherein G is hydrogen and R<sup>1</sup> is -SO<sub>2</sub>NR<sub>4</sub>R<sub>5</sub>. Referring to scheme 4, a compound of formula X wherein R<sup>1</sup> is -SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup> and G is (C<sub>1</sub>-C<sub>3</sub>)alkoxy is reacted with a reducing agent in a reaction inert solvent, for example  
15 lithium borohydride (LiBH<sub>4</sub>) in tetrahydrofuran (THF). The reduction, which yields an alcohol of the formula XVIII, is usually conducted at a temperature from about 0°C to about 100°C, preferably by heating the reaction mixture to the reflux temperature of the solvent. The alcohol of formula  
20 XVIII may then be oxidized using methods known to those skilled in the art. For example, treatment of a solution of such alcohol in a solvent such as methylene chloride with an oxidizing agent such as pyridinium dichromate at a temperature from about 0°C to about 50°C, preferably at room  
25 temperature, will yield the corresponding compounds of formula X wherein G is hydrogen and R<sup>1</sup> is -SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>. Other oxidizing agents/solvent systems such as manganese dioxide/acetone and chromium trioxide/acetic anhydride/acetic acid are also capable of producing this  
30 conversion.

The preparation of other compounds of the formula I not specifically described in the foregoing experimental section can be accomplished using combinations of the reactions described above that will be apparent to those skilled in  
35 the art.

In each of the reactions discussed or illustrated in schemes 1 to 4 above, pressure is not critical unless

-33-

otherwise indicated. Pressures from about 0.5 atmospheres to about 5 atmospheres are generally acceptable, and ambient pressure, i.e. about 1 atmosphere, is preferred as a matter of convenience.

5       The novel compounds of the formula I and the pharmaceutically acceptable salts thereof are useful as substance P antagonists, i.e., they possess the ability to antagonize the effects of substance P at its receptor site in mammals, and therefore they are able to function as  
10 therapeutic agents in the treatment of the aforementioned disorders and diseases in an afflicted mammal.

      The compounds of the formula I which are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although  
15 such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate a compound of the Formula I from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base  
20 compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a  
25 substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is readily obtained.

30       Those compounds of the formula I which are also acidic in nature, e.g., where R<sup>6</sup> or R<sup>10</sup> is carboxyphenyl, are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particularly,  
35 the sodium and potassium salts. These salts are all prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically

-34-

acceptable base salts of this invention are those which form non-toxic base salts with the acidic compounds of formula I. Such non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium, calcium and magnesium, etc. These salts can easily be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations, and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed in order to ensure completeness of reaction and maximum yields of the desired final product.

The compounds of formula I and their pharmaceutically acceptable salts exhibit substance P receptor-binding activity and therefore are of value in the treatment and prevention of a wide variety of clinical conditions the treatment or prevention of which are effected or facilitated by a decrease in substance P mediated neurotransmission. Such conditions include inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fasciitis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus

-35-

erythematosus, and rheumatic diseases such as fibrositis. Hence, these compounds are readily adapted to therapeutic use as substance P antagonists for the control and/or treatment of any of the aforesaid clinical conditions in mammals, including humans.

The compounds of the formula I and the pharmaceutically acceptable salts thereof can be administered via either the oral, parenteral or topical routes. In general, these compounds are most desirably administered in dosages ranging from about 5.0 mg up to about 1500 mg per day, although variations will necessarily occur depending upon the weight and condition of the subject being treated and the particular route of administration chosen. However, a dosage level that is in the range of about 0.07 mg to about 21 mg per kg of body weight per day is most desirably employed. Variations may nevertheless occur depending upon the species of animal being treated and its individual response to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval at which such administration is carried out. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into several small doses for administration throughout the day.

The compounds of the formula I and their pharmaceutically acceptable salts ("the therapeutic compounds") may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the three routes previously indicated, and such administration may be carried out in single or multiple doses. More particularly, the novel therapeutic agents of this invention can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies,

-36-

powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous  
5 media and various non-toxic organic solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the therapeutically-effective compounds of this invention are present in such dosage forms at concentration levels ranging  
10 from about 5.0% to about 70% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as  
15 starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are  
20 often very useful for tabletting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous  
25 suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water,  
30 ethanol, propylene glycol, glycerin and various like combinations thereof.

For parenteral administration, solutions of a therapeutic compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be  
35 employed. The aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous

-37-

injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

The activity of the therapeutic compounds of the present invention as substance P receptor antagonists may be determined by their ability to inhibit the binding of substance P at its receptor sites in bovine caudate tissue, employing radioactive ligands to visualize the tachykinin receptors by means of autoradiography. The substance P antagonizing activity of the herein described compounds may be evaluated by using the standard assay procedure described by M. A. Cascieri et al., as reported in the Journal of Biological Chemistry, Vol. 258, p. 5158 (1983). This method essentially involves determining the concentration of the individual compound required to reduce by 50% the amount of radiolabelled substance P ligands at their receptor sites in said isolated cow tissues, thereby affording characteristic  $IC_{50}$  values for each compound tested.

In this procedure, bovine caudate tissue is removed from a  $-70^{\circ}C$  freezer and homogenized in 50 volumes (w./v.) of an ice-cold 50 mM Tris (i.e., trimethamine which is 2-amino-2-hydroxymethyl-1,3-propanediol) hydrochloride buffer having a pH of 7.7. The homogenate is centrifuged at 30,000 x G for a period of 20 minutes. The pellet is resuspended in 50 volumes of Tris buffer, rehomogenized and then recentrifuged at 30,000 x G for another twenty-minute period. The pellet is then resuspended in 40 volumes of ice-cold 50 mM Tris buffer (pH 7.7) containing 2 mM of



-38-

calcium chloride, 2 mM of magnesium chloride, 4  $\mu$ g/ml of bacitracin, 4  $\mu$ g/ml of leupeptin, 2  $\mu$ g of chymostatin and 200 g/ml of bovine serum albumin. This step completes the production of the tissue preparation.

5       The radioligand binding procedure is then carried out in the following manner, viz., by initiating the reaction via the addition of 100  $\mu$ l of the test compound made up to a concentration of 1  $\mu$ M, followed by the addition of  
10       100  $\mu$ l of radioactive ligand made up to a final concentration 0.5 mM and then finally by the addition of 800  $\mu$ l of the tissue preparation produced as described above. The final volume is thus 1.0 ml, and the reaction mixture is next vortexed and incubated at room temperature (ca. 20°C) for a period of 20 minutes. The tubes are then filtered  
15       using a cell harvester, and the glass fiber filters (Whatman GF/B) are washed four times with 50 mM of Tris buffer (pH 7.7), with the filters having previously been presoaked for a period of two hours prior to the filtering procedure. Radioactivity is then determined in a Beta counter at 53%  
20       counting efficiency, and the  $IC_{50}$  values are calculated by using standard statistical methods.

The ability of the therapeutic compounds of this invention to inhibit substance P induced effects in vivo may be determined by the following procedures "a" through "d".  
25       (Procedures "a" through "c" are described in Nagahisa et al., European Journal of Pharmacology, 217, 191-5 (1992), which is incorporated herein by reference in its entirety.)

a. Plasma extravasation in the skin

Plasma extravasation is induced by intradermal  
30       administration of substance P (50  $\mu$ l, 0.01% BSA-saline solution) in dorsal skin of pentobarbital (25 mg/kg i.p.) anesthetized male Hartley guinea pigs weighing 450-500 g. The compound to be tested is dissolved in 0.1% methyl cellulose-water (MC) and dosed p.o. 1 hour before substance  
35       P challenge (3 pmol/site). Evans blue dye (30 mg/kg) is administered intravenously 5 minutes before challenge. After 10 minutes, the animals are sacrificed, the dorsal

-39-

skin is removed, and the blue spots are punched out using a cork borer (11.5 mm oral dose (o.d.)). Tissue dye content is quantitated after overnight formamide extraction at 600 nm absorbance.

5        b.    Capsaicin-induced plasma extravasation

Plasma extravasation is induced by intraperitoneal injection of capsaicin (10 ml of 30  $\mu$ M solution in 0.1% BSA/saline) into pentobarbital anesthetized (25 mg/kg i.p.) guinea pigs. The compound to be tested is dissolved in 0.1% MC and dosed p.o. 1 hour before capsaicin challenge. Evans blue dye (30 mg/kg) is administered i.v. 5 minutes before challenge. After 10 minutes, the animals are sacrificed, and both right and left ureters are removed. Tissue dye content is quantitated as in "a" above.

15       c.    Acetic acid-induced abdominal stretching

Male ddY mice (SLC, Japan), weighing 14-18 g, were fasted overnight. The compound to be tested is dissolved in 0.1% MC and dosed p.o. 0.5 hour before acetic acid (AA) injection (0.7%, 0.16 ml/10 g body weight). The animals are placed in clear beakers (1 per beaker) and the stretching response is counted 10 to 20 minutes after the AA injection (10 minute interval).

20       d.    Substance P-induced hyperlocomotor paradigm

The anti-psychotic activity of the therapeutic compounds of the present invention as neuroleptic agents for the control of various psychotic disorders may be determined by a study of their ability to suppress substance P-induced or substance P agonist induced hypermotility in guinea pigs. This study is carried out by first dosing the guinea pigs with a control compound or with an appropriate test compound of the present invention, then injecting the guinea pigs with substance P or a substance P agonist by intracerebral administration via canula and thereafter measuring their individual locomotor response to said stimulus.

35       The present invention is illustrated by the following examples. It will be understood, however, that the invention is not limited to the specific details of these examples.

-40-

PREPARATION 12-Cyclopentyloxy-5-(N-methyl-N-methanesulfonylamino)-benzaldehydeA. 4-Cyclopentyloxy-N-methanesulfonylaniline

5 Under N<sub>2</sub> in a flame-dried flask, a mixture of 4-cyclopentyloxyaniline (1.0 g, 5.64 mmol) in 25 mL of dry CH<sub>2</sub>Cl<sub>2</sub> was treated with triethylamine (1.3 mL, 9.38 mmol) and cooled to 0°C. A solution of recrystallized methanesulfonic anhydride (1.5 g, 8.62 mmol) in 10 mL of dry CH<sub>2</sub>Cl<sub>2</sub> was added  
10 dropwise and the reaction was stirred for 1.5 hours. The reaction mixture was then poured into 100 mL of saturated aqueous NaHCO<sub>3</sub> and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 X 50 mL). The combined organics were dried over MgSO<sub>4</sub>, evaporated in vacuo to a dark grey solid and flash chromatographed on silica  
15 gel, eluting with hexanes:EtOAc (80:20), to produce the pure intermediate compound, 0.8 g (56%), m.p. 140-142°C.

B. 4-Cyclopentyloxy-N-methyl-N-methanesulfonylaniline

Under N<sub>2</sub> in a flame-dried flask, the preceding intermediate (0.5 g, 1.96 mmol) in 25 mL of acetone was  
20 treated with K<sub>2</sub>CO<sub>3</sub> (0.54 g, 3.91 mmol), stirred for 5 minutes at 25°C and treated with methyl iodide (0.33 g, 2.32 mmol). After 18 hours, the suspension was filtered through a pad of diatomaceous earth (d.e.), concentrated in vacuo, redissolved in 100 mL of ethyl acetate (EtOAc), refiltered and  
25 concentrated to an off-white solid, 300 mg (57%), m.p. 120-122°C.

C. 2-Cyclopentyloxy-5-(N-methyl-N-methanesulfonylamino)benzaldehyde

Under N<sub>2</sub> in a flame-dried flask, the above intermediate  
30 from part "B" (300 mg, 1.11 mmol) in 15 mL of CH<sub>2</sub>Cl<sub>2</sub> was cooled to 0°C and treated with titanium tetrachloride (0.46 g, 0.27 mL, 2.42 mmol). After 20 minutes at 0°C α,α-dichloromethyl methyl ether (0.15 g, 0.12 mL, 1.33 mmol) was added and the reaction was left to slowly warm to room  
35 temperature overnight. The reaction was quenched in 100 mL of saturated aqueous NaHCO<sub>3</sub>, extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 X 75 mL) and dried over MgSO<sub>4</sub>. Concentration in vacuo gave a light

-41-

brown solid which was filtered through a pad of d.e. to obtain the purified title aldehyde, 155 mg (47%).

Mass Spectrum (MS): m/e 297 (p+), 229, 150 (100%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.6-2.05 (m, 10H), 2.85 (s, 3H), 3.3  
5 (s, 3H), 4.95 (m, 1H), 7.0 (d, 1H), 7.2-7.75 (m, 3H), 10.5 (s, 1H).

The following intermediate aldehydes of the general formula X were prepared by a procedure similar to that of Preparation 1.

10 2-Methoxy-5-(trifluoromethylthio)benzaldehyde, m.p. 61-64°C, 30% yield.

5-Tert-butyl-2-methylthiobenzaldehyde, oil, 54% yield, MS: m/e 208 (p+), 193 (100%), 165, 117.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.30 (s, 9H), 2.45 (s, 3H), 7.28 (d,  
15 1H), 7.55 (dd, 1H), 7.82 (d, 1H), 10.3 (s, 1H).

5-Chloro-2-methylthiobenzaldehyde, m.p. 51-54°C, 52% yield.

2-Methoxy-5-(N-methyl-N-methanesulfonylamino)-benzaldehyde, 89%, <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 2.9 (s, 3H), 3.4 (s, 3H),  
20 4.0 (s, 3H), 7.1 (d, 1H), 7.7-7.85 (m, 2H), 10.5 (s, 1H).

2-Methoxy-5-(N-isopropyl-N-methanesulfonylamino)-benzaldehyde, m.p. 114-116°C, 81% yield.

2-Methoxy-5-(1,1-dioxo-2-thiazolidinyl)benzaldehyde, m.p. 99-101°C, 82% yield.

25 2-Isopropoxy-5-(N-methyl-N-methanesulfonylamino)-benzaldehyde, m.p. 107-110°C, 60% yield.

2-Isopropoxy-5-(N-methyl-N-trifluoromethanesulfonylamino)benzaldehyde, m.p. 42-45°C, 83% yield.

2-Methoxy-5-(N-methyl-N-trifluoromethanesulfonylamino)-benzaldehyde, 55% yield, <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.46 (s,  
30 3H), 3.98 (s, 3H), 7.04 (d, 1H), 7.57 (dd, 1H), 7.81 (d, 1H), 10.4 (s, 1H). MS: m/e 297 (p+).

2-Methoxy-5-(N-methyl-N-isopropylsulfonylamino)-benzaldehyde, 39% yield.

35 2-Methoxy-5-(N-methyl-N-(4-methylphenylsulfonyl)-amino)-benzaldehyde, oil, 88% yield.

-42-

2-Isopropoxy-5-(N-methyl-N-(4-methylphenylsulfonyl)-amino)benzaldehyde, 28% yield,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.45 (d, 6H), 2.42 (s, 3H), 3.10 (s, 3H), 4.70 (m, 1H), 7.0 (m, 3H), 7.25 (m, 3H), 7.42 (d, 2H), 7.58 (dd, 1H), 10.4 (s, 1H). MS: m/e 347 (p+), 305, 150 (100%).

2-Methoxy-5-(N-methyl-N-benzylsulfonylamino)-benzaldehyde, 51% yield,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.14 (s, 3H), 3.94 (s, 3H), 4.27 (s, 2H), 6.95 (d, 1H), 7.35-7.58 (m, 7H), 10.4 (s, 1H). MS: m/e 319 (p+), 255, 164, 91 (100%).

10 5-Methoxy-1-methanesulfonyl-2,3-dihydroindol-6-carboxaldehyde, 49% yield,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.85 (s, 3H), 3.19 (t, 2H), 3.90 (s, 3H), 3.98 (t, 2H), 6.90 (s, 1H), 7.73 (s, 1H), 10.3 (s, 1H).

5-Methoxy-3-methyl-1-methanesulfonyl-2,3-dihydroindol-15 6-carboxaldehyde, m.p. 147-150°C, 49% yield,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.45 (d, 3H), 2.75 (dd, 1H), 2.85 (s, 3H), 3.5 (dd, 1H), 3.95 (s, 3H), 4.5 (m, 1H), 6.9 (s, 1H), 7.8 (s, 1H), 10.4 (s, 1H).

2-Methoxy-5-(N-cyclopentyl-N-(4-methanesulfonylamino)-20 benzaldehyde, m.p. 95-98°C, 62% yield.

2-Methoxy-5-(2-methyl-4-thiazolyl)benzaldehyde, 56% yield,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.72 (s, 3H), 3.95 (s, 3H), 7.05 (d, 1H), 7.25 (s, 1H), 8.15 (dd, 1H), 8.25 (d, 1H), 10.5 (s, 1H).

25 2-Methoxy-5-(N-(3,4-dichlorobenzyl)-N-methanesulfonylamino)benzaldehyde, gum, 86% yield,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.97 (s, 3H), 3.95 (s, 3H), 4.75 (s, 2H), 6.95 (d, 1H), 7.10 (d, 1H), 7.35 (m, 3H), 7.77 (d, 1H), 10.4 (s, 1H).

2-Methoxy-5-(N-cyclohexylmethyl-N-methanesulfonyl-30 amino)benzaldehyde, oil, 73% yield,  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.9-1.8 (m, 11H), 2.85 (s, 3H), 3.48 (d, 2H), 3.98 (s, 3H), 7.05 (d, 1H), 7.60 (dd, 1H), 7.75 (d, 1H), 10.5 (s, 1H).

5-(Isopropylsulfonyl)-2-methoxybenzaldehyde, m.p. 105-107°C, 57% yield, MS: m/e 242 ( $\text{M}^+$ , 27%), 200 (78%), 136 35 (100%),  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.3 (d, 6H), 3.15 (m, 1H), 4.05 (s, 3H), 7.15 (d, 1H), 8.05 (dd, 1H), 8.3 (d, 1H), 10.5 (s, 1H).

-43-

5-(N-cyclopentyl-N-methanesulfonyl)amino-2-methoxybenzaldehyde, m.p. 95-98°C, 62% yield, MS: m/e 297 ( $M^+$ , 20%), 229, 218 (100%), 150 (95%),  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.25-1.6 (m, 6H), 1.95 (m, 2H), 2.95 (s, 3H), 3.95 (s, 3H), 4.5 (m, 1H), 7.05 (d, 1H), 7.5 (dd, 1H), 7.7 (d, 1H), 10.45 (s, 1).

5-(N-cyclohexylmethyl-N-methanesulfonyl)amino-2-methoxybenzaldehyde, oil, 74% yield,  $^1H$  NMR ( $CDCl_3$ )  $\delta$  0.9-1.8 (m, 11H), 2.85 (s, 3H), 3.45 (d, 2H), 4.0 (s, 3H), 7.05 (d, 1H), 7.65 (dd, 1H), 7.75 (d, 1H), 10.45 (s, 1H).

2,3-Dihydro-N-methanesulfonyl-5-methoxy-2-methylindole-6-carboxaldehyde, m.p. 147-150°C, 48% yield,  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.45 (d, 3H), 2.75 (m, 1H), 2.85 (s, 3H), 3.5 (dd, 1H), 3.9 (s, 3H), 4.5 (m, 1H), 6.9 (s, 1H), 7.83 (s, 1H), 10.4 (s, 1H).

2-Methoxy-5-(N-methyl-N-(2,4-dimethyl-5-thiazolesulfonyl))aminobenzaldehyde, oil, 29% yield, MS: m/e 340 ( $M^+$ , 10%), 164 (100%),  $^1H$  NMR ( $CDCl_3$ )  $\delta$  2.1 (s, 3H), 2.5 (s, 3H), 3.1 (s, 3H), 3.9 (s, 3H), 7.0 (d, 1H), 7.5 (m, 1H), 7.6 (q, 1H), 10.4 (s, 1H).

2-Methoxy-5-(N-(4,5-dimethyl-2-thiazolyl)-N-methanesulfonyl)aminobenzaldehyde, waxy solid, 39% yield, MS: m/e (340 ( $M^+$ , 20%), 261 (65%),  $^1H$  NMR ( $CDCl_3$ )  $\delta$  2.3 (d, 6H), 3.4 (s, 3H), 4.0 (s, 3H), 7.0 (s, 3H), 7.0 (d, 1H), 7.7 (q, 1H), 10.5 (s, 1H).

2-Methoxy-5-(N-(4,5-dimethyl-2-thiazolyl)-N-methyl)aminobenzaldehyde, oil, 7% yield, MS: m/e 277 ( $M^+$ , 20%), 276 (100%), 126 (30),  $^1H$  NMR ( $CDCl_3$ )  $\delta$  2.1 (d, 6H), 3.4 (s, 3H), 4.0 (s, 3H), 7.0 (d, 1H), 7.6 (q, 1H), 7.8 (d, 1H), 10.5 (s, 1H).

2-Methoxy-5-(N-(4,5-dimethyl-2-thiazolyl))aminobenzaldehyde, m.p. 137-139°C, 20% yield, MS: m/e 262 ( $M^+$ , 100%),  $^1H$  NMR ( $CDCl_3$ )  $\delta$  2.15 (s, 3H), 2.25 (s, 3H), 3.9 (s, 3H), 7.0 (d, 1H), 7.6 (dd, 1H), 7.7 (dd, 1H), 10.5 (s, 1H).

2-Methoxy-5-(N-(ethoxycarbonylmethanesulfonyl)-N-methyl)aminobenzaldehyde, oil, 81% yield,  $^1H$  NMR ( $CDCl_3$ )  $\delta$

-44-

1.3 (t, 3H), 2.05 (s, 2H), 3.35 (s, 3H), 3.9 (s, 3H), 4.25 (q, 2H), 7.05 (d, 1H), 7.7 (dd, 1H), 7.9 (d, 1H), 10.5 (s, 1H).

2-Methoxy-5-(N-(3,4-dichlorobenzyl)-N-methanesulfonyl)aminobenzaldehyde, 86% yield, <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.0 (s, 3H), 3.95 (s, 3H), 4.8 (s, 2H), 6.95 (d, 1H), 7.15 (dd, 1H), 7.35 (m, 3H), 7.75 (d, 1H), 10.4 (s, 1H).

#### PREPARATION 2

##### 2-Methoxy-5-methylthiobenzaldehyde

Under N<sub>2</sub> in a flame-dried flask, fitted with a condensor and stirrer, was placed a solution of 1-methoxy-4-methylthiobenzene (2.0 g, 13 mmol) in 75 mL of trifluoroacetic acid (TFA). Hexamethylenetetramine (1.2 g, 13 mmol) was added while stirring the reaction at 25°C. After heating for 2 hours to reflux, the reaction was cooled and concentrated in vacuo and the residue was partitioned between CH<sub>2</sub>Cl<sub>2</sub> and 2 N sodium hydroxide (NaOH). The organic layer was dried over MgSO<sub>4</sub>, concentrated in vacuo to a yellow oil and flash chromatographed on silica gel eluting with hexanes:EtOAc (85:15) to give the pure title compound as a yellow oil, 0.99 g, 42% yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 2.5 (s, 3H), 3.95 (s, 3H), 6.97 (d, 1H), 7.5 (dd, 1H), 7.78 (d, 1H), 10.5 (s, 1H).

Using a procedure similar to that of Preparation 2, 4-methoxyphenyl cyclohexyl sulfide was converted to 2-methoxy-5-(cyclohexylthio)benzaldehyde, oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.1-2.0 (m, 10H), 3.0 (m, 1H), 3.95 (s, 3H), 6.95 (d, 1H), 7.62 (dd, 1H), 7.90 (d, 1H), 10.45 (s, 1H).

#### PREPARATION 3

##### 2-Phenoxybenzaldehyde

A mixture of 2-phenoxybenzyl alcohol (4.0 g, 20.2 mmol, prepared by the reduction of commercially available 2-phenoxybenzoic acid with LiAlH<sub>4</sub>/THF) and 150 mL CH<sub>2</sub>Cl<sub>2</sub> was treated with pyridinium dichromate (11.39 g, 30.3 mmol) at 25°C and stirred for another 36 hours. The mixture was filtered through d.e. and then through a pad of silica gel to produce 3.11 g (78%) of title compound as a yellow oil.

-45-

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 6.8 (d, 1H), 6.9 (d, 1H), 7.0 (m, 2H), 7.1 (m, 2H), 7.2 (m, 1H), 8.9 (m, 1H), 10.2 (s, 1H).

Using a procedure similar to that of Preparation 3, the following benzaldehydes of formula X were prepared from the  
5 corresponding compounds of formula XVIII:

5-Diethylaminosulfonyl-2-methoxybenzaldehyde, 76%, <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.1 (t, 6H), 3.2 (q, 4H), 4.0 (s, 3H), 7.1 (d, 1H), 7.9 (dd, 1H), 8.2 (d, 1H), 10.4 (s, 1H).

5-Diethylaminosulfonyl-2-isopropoxybenzaldehyde, oil,  
10 36%, <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.2 (t, 6H), 1.4 (d, 6H), 3.2 (q, 4H), 4.8 (m, 1H), 7.1 (d, 1H), 8.0 (dd, 1H), 8.2 (d, 1H), 10.4 (s, 1H).

#### PREPARATION 4

##### 2-Trifluoromethoxy-5-(N-methyl-N-methanesulfonyl)- 15 benzaldehyde

A. To a mixture of concentrated sulfuric acid (81 mL) and concentrated nitric acid (15.5 mL), cooled to 0°C, was added 2-trifluoromethoxybenzaldehyde (25 g, 0.13 mol) portionwise while maintaining the temperature of the  
20 reaction below 0°C. After 1.5 hours, the reaction mixture was poured cautiously over 1000 mL of ice in a large beaker and left to stand for 0.5 hours. The resulting suspension was filtered, washed well with H<sub>2</sub>O and air dried to give crude 5-nitro-2-trifluoromethoxybenzaldehyde, m.p. 32-34°C.

25 B. The preceding compound and ethylene glycol (35 mL, 0.62 mol) in 1000 mL of toluene was treated with para-toluenesulfonic acid (0.72 g, 4 mmol) and heated to reflux under N<sub>2</sub> for 24 hours, using a Dean-Stark trap to collect the water formed. The solvent was then removed in vacuo and the  
30 residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, washed with saturated aqueous NaHCO<sub>3</sub> and dried over MgSO<sub>4</sub>. Removal of the solvent in vacuo gave crude 2-(5-nitro-2-trifluoromethoxyphenyl)-1,3-dioxolane as a pale orange oil.

C. The dioxolane from part B (5.09 g) in 100 mL of  
35 EtOAc was hydrogenated with 0.29 g of 5% palladium on carbon at 45 p.s.i. for 18 hours. After filtration through d.e., the solvent was removed in vacuo to give 2-(5-amino-2-



-46-

trifluoromethoxyphenyl)-1,3-dioxolane as an orange oil, 4.6 g.

D. The above oil from part C and triethylamine (2.53 mL, 39 mmol) in 200 mL of dry THF was treated with  
5 methanesulfonic anhydride (4.9 g, 28 mmol) in 26 mL of THF at 25°C. After 72 hours, 200 mL of H<sub>2</sub>O was added and the mixture was stirred for another 30 minutes. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the organics were combined, washed with 1 N HCl, 2 N NaOH and H<sub>2</sub>O, and finally  
10 dried over MgSO<sub>4</sub>. Removal of the solvent in vacuo gave an orange oil which was flash chromatographed on silica gel using hexanes:EtOAc (40:60). Pure 2-(5-methanesulfonylamino-2-trifluoromethoxyphenyl)-1,3-dioxolane was obtained as an oil, 1.64 g.

15 E. A suspension of sodium hydride (60% oil dispersion, 0.19 g, 4.75 mmol) in 10 mL of dry DMF was treated with the compound from part D (1.5 g, 4.58 mmol) in 20 mL of dry DMF and stirred at 25°C for 30 minutes. Methyl iodide (0.28 mL, 4.5 mmol) was added and the mixture was  
20 stirred for an additional 15 hours. After dilution with 100 mL of water, the mixture was extracted with Et<sub>2</sub>O (3 X 100 mL) and the combined organics were dried over MgSO<sub>4</sub> and evaporated to give 2-(5-N-methyl-N-methanesulfonylamino)-2-trifluoromethoxyphenyl)-1,3-dioxolane as an orange oil,  
25 1.63 g.

F. The preceding dioxolane from part E (1.63 g) in 30 mL of acetone was treated with 6 N HCl at 25°C for 72 hours. The acetone was then evaporated and the resulting solution was extracted with Et<sub>2</sub>O and the organics were washed with  
30 H<sub>2</sub>O, dried over MgSO<sub>4</sub> and concentrated to an oil. Flash chromatography on silica gel using hexanes:EtOAc (65:35) gave pure 5-(N-methyl-N-methanesulfonylamino)-2-trifluoromethoxy-benzaldehyde as an oil, 0.63 g (42%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 2.9 (s, 3H), 3.4 (s, 3H), 7.4 (dd, 1H), 7.8  
35 (dd, 1H), 7.9 (d, 1H), 10.4 (s, 1H). MS: m/e 297 (p+), 218, 162.

-47-

PREPARATION 52-Methoxy-5-methanesulfonylbenzaldehyde

Under N<sub>2</sub> in a round-bottomed flask fitted with a condensor, 2-methoxy-5-methylthiobenzaldehyde (0.89 g, 4.9 mmol) was added to 0.6 mL of EtOH. To this, a solution of monoperoxyphthalic acid magnesium salt hexahydrate (2.41 g, 4.9 mmol) in 10.4 mL of H<sub>2</sub>O was added and the mixture was heated at 95°C for 18 hours. The reaction was then quenched with 10 mL of H<sub>2</sub>O, extracted with CH<sub>2</sub>Cl<sub>2</sub> (4 X 10 mL) and the combined organics were dried over MgSO<sub>4</sub> and concentrated in vacuo to an oil, 0.34 g. Flash chromatography on silica gel, eluting with EtOAc:hexanes (2.98) gave the pure title compound as a white solid, 0.36 g, m.p. 140-143°C. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 3.05 (s, 3H), 4.05 (s, 3H), 7.2 (d, 1H), 8.15 (dd, 1H), 8.40 (d, 1H), 10.5 (s, 1H).

EXAMPLE 1Cis-3-(5-fluoro-2-methylthiobenzyl)amino-2-phenylpiperidine dihydrochlorideA. 5-Fluoro-2-methylthiobenzaldehyde

Under N<sub>2</sub> a solution of 8.81 g (62 mmole) of p-fluorothioanisole in 50 mL of dichloromethane was stirred, cooled to 0°C and treated dropwise with 15 mL (136 mmole) of titanium tetrachloride (TiCl<sub>4</sub>). After stirring approximately 30 minutes at this temperature, the red solution was treated with 6.73 mL (74.4 mmole) of a α,α-dichloromethyl methyl ether (Aldrich Chem Co.), stirred an additional 2 hours at 0°C and allowed to warm to room temperature while stirring for another 18 hours. After pouring the reaction mixture into a mixture of 250 mL of saturated aqueous sodium bicarbonate and 250 mL of dichloromethane, the aqueous layer was extracted with three 50 mL portions of dichloromethane and the organic layers were combined and dried over magnesium sulfate (MgSO<sub>4</sub>). Evaporation of the solvent produced a solid which was recrystallized from hexane; 0.72 g, M.P. 45-47°C. Mass spectrum (m/e, %); 172 (17), 171 (33), 170 (100, M<sup>+</sup>), 155 (49), 142 (53), 127 (28).

B. Cis-3-(5-fluoro-2-methylthiobenzyl)amino-6-oxo-2-phenylpiperidine

A mixture of 0.67 g (3.52 mmole) of cis-3-amino-6-oxo-2-phenylpiperidine, 0.72 g (4.23 mmole) of the above  
5 aldehyde and 1 g of 3A molecular sieves (Aldrich) in 15 mL of acetic acid was stirred at 25°C for approximately 1.5 hours, then treated with 1.71 g (8.1 mmole) of sodium triacetoxyborohydride. After stirring for another 18 hours, the mixture was filtered and the filtrate concentrated to a  
10 yellow oil. Chromatography on silica gel using dichloromethane:ethanol:concentrated ammonium hydroxide (98:1:1) produced the pure product as an oil which crystallized on standing; 0.51 g (42%), M.P. 125-130°C. Mass spectrum (m/e, %): 345 (45, M<sup>+</sup>) 344 (100 M<sup>+</sup>), 210  
15 (92), 155 (91), 106 (99).

C. Cis-3-(5-fluoro-2-methylthiobenzyl)amino-2-phenylpiperidine dihydrochloride

In a flame-dried flask 0.69 g (2 mmole) of the previous compound in 5 mL of tetrahydrofuran was treated with 3.0 mL  
20 of 1.0 M borane-tetrahydrofuran complex (Aldrich), refluxed for 1 hour and stirred at 25°C for 18 hours. After acidifying the crude mixture with 2N hydrochloric acid (HCl), it was extracted with dichloromethane and the aqueous layer was made basic with 2N sodium hydroxide (NaOH). This  
25 alkaline layer was finally extracted with dichloromethane which was dried over (MgSO<sub>4</sub>) and concentrated to an oil; on standing it crystallized to an off-white solid, mp 60-64°C. This free base was redissolved in dichloromethane and treated with hydrogen chloride (HCl) gas to form the  
30 dihydrochloride salt, recrystallized from methanol:diethyl ether as a white crystalline solid, M.P. 270-273°C. Mass spectrum (m/e, %): 330 (15, M<sup>+</sup>), 211 (100), 210 (65), 155 (98).

Anal. calc'd for C<sub>19</sub>H<sub>23</sub>FN<sub>2</sub>S•0.5 H<sub>2</sub>O: C, 55.34; H, 6.35;  
35 N, 6.79. Found: C, 55.08; H, 6.51; N, 6.59.

The title compounds of Examples 2-8 were prepared by a procedure similar to that of Example 1.

-49-

EXAMPLE 2Cis-3-(2-methylthiophenyl)methylamino-2-phenylpiperidine dihydrochlorideM.P. 256-259°C (MeOH:Et<sub>2</sub>O)5 MS (m/e, %): 312 (M<sup>+</sup>), 193, 192, 175, 160, 137 (100)Anal. calc'd for C<sub>19</sub>H<sub>24</sub>N<sub>2</sub>S•2HCl: C, 59.21; H, 6.80; N, 7.27. Found: C, 59.08; H, 6.92; N, 7.18EXAMPLE 310 Cis-3-(5-tert-butyl-2-methylthiophenyl)methylamino-2-phenylpiperidine dihydrochlorideM.P. 237-240°C (MeOH:Et<sub>2</sub>O)MS (m/e, %): 368 (3, M<sup>+</sup>), 367, 264, 210, 175, 155.Anal. calc'd for C<sub>23</sub>H<sub>32</sub>N<sub>2</sub>S•2HCl•0.5 CH<sub>2</sub>Cl<sub>2</sub>: C, 59.65; H, 7.44; N, 6.79. Found: C, 59.37; H, 7.38; N, 6.12.

15

EXAMPLE 4Cis-3-(5-chloro-2-methylthiophenyl)methylamino-2-phenylpiperidine dihydrochlorideM.P. 260-265°C (MeOH:Et<sub>2</sub>O)MS (m/e, %): 348, 346 (M<sup>+</sup>), 227, 180, 171, 120, 106.20 Anal. calc'd for C<sub>19</sub>H<sub>23</sub>ClN<sub>2</sub>S•2HCl: C, 54.35; H, 6.00; N, 6.67. Found: C, 54.04; H, 6.08; N, 6.66.EXAMPLE 5Cis-3-(2-tert-butylthiophenyl)methylamino-2-phenylpiperidine dihydrochloride25 M.P. 243-245°C dec. (MeOH:Et<sub>2</sub>O)MS (m/e, %): 354 (6, M<sup>+</sup>), 297, 235, 234, 178, 160, 123, 70(100).Anal. calc'd for C<sub>22</sub>H<sub>30</sub>N<sub>2</sub>S•2HCl: C, 61.81; H, 7.55; N, 6.55. Found: C, 61.46; H, 7.26; N, 6.52.

30

EXAMPLE 6Cis-3-(2-(4-chlorophenylthio)phenyl)methylamino-2-phenylpiperidine dihydrochlorideM.P. 245-249°C dec. (MeOH:Et<sub>2</sub>O)35 MS (m/e, %): 408 (M<sup>+</sup>), 289, 231, 197 (100), 165, 146, 120.Anal. calc'd for C<sub>24</sub>H<sub>25</sub>ClN<sub>2</sub>S•2HCl•1/3 H<sub>2</sub>O: C, 59.08; H, 5.72; N, 5.74. Found: C, 59.08; H, 5.61; N, 5.84.

-50-

EXAMPLE 7Cis-3-(2-methoxy-5-(trifluoromethylthio)phenyl)methyl-amino-2-phenylpiperidine dihydrochlorideM.P. 257-259°C (MeOH:Et<sub>2</sub>O)

5        Anal. calc'd for C<sub>20</sub>H<sub>23</sub>F<sub>3</sub>N<sub>2</sub>OS•2HCl•1/2H<sub>2</sub>O: C, 50.21; H, 5.48; N, 5.86. Found: C, 50.60; H, 5.42; N, 6.09.

EXAMPLE 8Cis-3-(2-phenoxyphenyl)methylamino-2-phenylpiperidine hydrochloride10        M.P. 210-212°C (MeOH:Et<sub>2</sub>O)

MS (m/e, %): 358 (M<sup>+</sup>), 239, 198, 183 (100), 175, 160, 146.

Anal. calc'd for C<sub>24</sub>H<sub>26</sub>N<sub>2</sub>O•HCl•1/4H<sub>2</sub>O; C, 71.97; H, 6.79; N, 6.91. Found: C, 72.16; H, 6.94; N, 7.01.

15

EXAMPLE 9(+)-(2S,3S)-3-[2-methoxy-5-(N-isopropyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine dihydrochlorideA. (+)-(2S,3S)-3-Amino-2-phenylpiperidine

20        In a bottle were placed 9 g of 10 % palladium-carbon, 180 ml of methanol, 275 ml of ethanol, 6.5 ml of concentrated hydrochloric acid and 9 g of the hydrochloride salt of (2S,3S)-3-(2-methoxybenzylamino)-2-phenylpiperidine. The mixture was shaken under hydrogen (40 p.s.i.) overnight, 25 after which 9 g of additional catalyst were added to the system and the mixture was shaken under hydrogen for 1 day. The mixture was diluted with water (250 mL), filtered through diatomaceous earth (Celite (trademark)) and the Celite was rinsed well with water. The filtrate was 30 concentrated to a volume of ca. 600-700 mL, made basic with concentrated aqueous sodium hydroxide and extracted with chloroform. The chloroform extracts were dried (sodium sulfate) and concentrated to obtain 4.4 g of the title compound as a colorless oil.

35        [α]<sub>D</sub> (HCl salt) = + 62.8° (c = 0.46, methanol (CH<sub>3</sub>OH)).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.68 (m, 4H), 2.72 (m, 1H), 2.94 (broad s, 1H), 3.16 (m, 1H), 3.80 (d, 1H, J=3), 7.24 (m, 5H).

-51-

HRMS calc'd for  $C_{11}H_{16}N_2$ : 176.1310. Found: 176.1309.

Anal. calc'd for  $C_{11}H_{16}N_2 \cdot 2HCl \cdot 1/3H_2O$ : C, 51.78; H, 7.36; N, 10.98. Found: C, 51.46; H, 7.27; N, 10.77.

B. (+)-(2S,3S)-3-[2-methoxy-5-(N-isopropyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine dihydrochloride

Under a nitrogen atmosphere in a round-bottom flask were placed 80 mg (0.46 mmol) of (+)-(2S,3S)-3-amino-2-phenylpiperidine, 5 ml of acetic acid and 150 mg (0.55 mmol) of 2-methoxy-5-(N-isopropyl-N-methanesulfonylamino)benzaldehyde, and the mixture was stirred for 60 minutes. To the system were added 0.21 g (1.0 mmol) of sodium triacetoxyborohydride, and the mixture was stirred at room temperature overnight. The mixture was concentrated, basified with 1 M aqueous sodium hydroxide and extracted with methylene chloride. The methylene chloride extracts were washed with water and extracted with 1 M aqueous hydrochloric acid. The hydrochloric acid extracts were basified with 1 M aqueous sodium hydroxide and extracted with methylene chloride. The methylene chloride extracts were dried (sodium sulfate) and concentrated to obtain 528 mg of colorless oil. The oil was dissolved in methylene chloride, and ether saturated with hydrogen chloride was added to the solution. The resulting white solid was collected by filtration and stirred in isopropanol at 60°C for 2 hours. Filtration afforded 128 mg of the title compound as its hydrochloride.

M.P. 268-270°C.

$^1H$  NMR ( $CDCl_3$ ; free base)  $\delta$  1.0 (d, 6H), 1.38-2.20 (m, 6H), 2.80 (m, 2H), 2.85 (s, 3H), 3.2 (t, 1H), 3.35 (d, 1H), 3.50 (s, 3H), 3.70 (d, 1H), 3.90 (d, 1H), 4.50 (m, 1H), 6.65 (d, 1H), 6.90 (d, 1H), 7.05 (dd, 1H), 7.25 (m, 5H).

Mass spectrum: m/z 431 (parent), 312 (100%).

Anal. calc'd for  $C_{27}H_{33}N_3O_3S \cdot 2HCl$ : C, 54.75; H, 6.99; N, 8.32. Found: C, 54.75; H, 6.99; N, 8.29.

The title compounds of Examples 10-37 were prepared from either (+)-(2S,3S)-3-amino-2-phenylpiperidine or the

-52-

corresponding racemate by employing the appropriate aldehyde and using a procedure similar to that of Example 9B.

EXAMPLE 10

(2S,3S)-3-(2-Methoxy-5-methylmercaptobenzylamino)-2-phenylpiperidine hydrochloride

M.P. 257 - 259°C (dec.)

<sup>1</sup>H NMR (free base; CDCl<sub>3</sub>) δ 1.32 (m, 1H), 1.50 (m, 1H), 1.82 (m, 1H), 2.04 (m, 1H), 2.30 (s, 3H), 2.72 (m, 2H), 3.18 (m, 1H), 3.26 (d, 1H, J=15), 3.36 (s, 3H), 3.54 (d, 1H, J=15), 3.80 (d, 1H, J=3), 6.52 (d, 1H, J=10), 6.90 (d, 1H, J=3), 7.04 (dd, 1H, J=3, 10), 7.2 (m, 5H).

HRMS calc'd for C<sub>20</sub>H<sub>26</sub>N<sub>2</sub>OS: 342.1760. Found: 342.1770.

Anal. calc'd for C<sub>20</sub>H<sub>26</sub>N<sub>2</sub>OS•2HCl•0.25H<sub>2</sub>O: C, 57.20; H, 6.84; N, 6.67. Found: C, 57.35; H, 6.76; N, 6.61.

15

EXAMPLE 11

(2S,3S)-3-(2-Methoxy-5-methylsulfoxybenzylamino)-2-phenylpiperidine hydrochloride

M.P. 209°C (dec).

<sup>1</sup>H NMR (free base; CDCl<sub>3</sub>) δ 1.40 (m, 1H), 1.56 (m, 1H), 1.90 (m, 1H), 2.10 (m, 1H), 2.59, 2.62 (2S, 3H), 2.76 (m, 2H), 3.22 (m, 1H), 3.42 (m, 1H), 3.49, 3.52 (2S, 3H), 3.66 (m, 1H), 3.86 (d, 1H, J=3), 6.76 (m, 1H), 7.24 (m, 6H), 7.46 (m, 1H).

HRMS calc'd for C<sub>20</sub>H<sub>27</sub>N<sub>2</sub>O<sub>2</sub>S(M+1): 359.1787. Found: 359.1763.

25

EXAMPLE 12

(2S,3S)-3-(2-Methoxy-5-methylsulfonylbenzylamino)-2-phenylpiperidine hydrochloride

M.P. > 260°C.

<sup>1</sup>H NMR (free base; CDCl<sub>3</sub>) δ 1.40 (m, 1H), 1.58 (m, 1H), 1.88 (m, 1H), 2.10 (m, 1H), 2.78 (m, 2H), 2.96 (s, 3H), 3.24 (m, 1H), 3.38 (d, 1H, J=15), 3.54 (s, 3H), 3.66 (d, 1H, J=15), 3.90 (d, 1H, J=3), 6.74 (d, 1H, J=10), 7.26 (m, 5H), 7.58 (d, 1H, J=3), 7.72 (d, 1H, J=10).

HRMS calc'd for C<sub>20</sub>H<sub>26</sub>N<sub>2</sub>O<sub>3</sub>S: 374.1658. Found: 374.1622.

35

-53-

EXAMPLE 13(2S,3S)-3-(2-Methoxy-5-phenoxybenzylamino)-2-phenylpiperidine hydrochloride

M.P. &gt; 250°C.

- 5       <sup>1</sup>H NMR (free base; CDCl<sub>3</sub>) δ 1.34 (m, 1H), 1.74 (m, 2H),  
2.06 (m, 1H), 2.76 (m, 2H), 3.22 (m, 1H), 3.32 (d, 1H,  
J=15), 3.44 (s, 3H), 3.60 (d, 1H, J=15), 3.85 (d, 1H, J=3),  
6.60 (d, 1H, J=9), 6.67 (d, 1H, J=3), 6.78 (dd, 1H, J=6,9),  
6.86 (d, 2H), 7.00 (t, 1H, J=6), 7.22 (m, 7H).
- 10       HRMS calc'd for C<sub>25</sub>H<sub>28</sub>N<sub>2</sub>O<sub>2</sub>: 388.2151. Found: 388.2137.

EXAMPLE 14(2S,3S)-3-(2-Methoxy-5-N-methylmethanesulfonylamino-benzylamino)-2-phenylpiperidine hydrochloride

M.P. 281-283°C.

- 15       <sup>1</sup>H NMR (free base; CDCl<sub>3</sub>) δ 1.42 (m, 1H), 1.74 (m, 2H),  
2.12 (m, 1H), 2.78 (m, 5H), 3.20 (s, 3H), 3.24 (m, 1H), 3.36  
(d, 1H, J=15), 3.52 (s, 3H), 3.64 (d, 1H, J=15), 3.89 (d,  
1H, J=3), 6.64 (d, 1H, J=9), 6.98 (d, 1H, J=3), 7.14 (dd,  
1H, J=3, 9), 7.26 (m, 5H).
- 20       HRMS calc'd for C<sub>21</sub>H<sub>29</sub>N<sub>3</sub>O<sub>3</sub>S: 403.1992. Found: 403.1923.  
Anal. calc'd for C<sub>21</sub>H<sub>29</sub>N<sub>3</sub>O<sub>3</sub>S•2HCl•1/3H<sub>2</sub>O: C, 52.28; H,  
6.61; N, 8.71. Found: C, 52.09; H, 6.63; N, 8.68.

EXAMPLE 15Cis-3-[2-isopropoxy-5-(N-methyl-N-methanesulfonyl-amino)benzylamino]-2-phenylpiperidine dihydrochloride

M.P. 278-280°C, 39% yield.

Anal. calc'd for C<sub>22</sub>H<sub>33</sub>N<sub>3</sub>O<sub>3</sub>S•2HCl: C, 54.75; H, 6.99; N,  
8.32. Found: C, 54.83, H, 7.16, N, 8.16.

- 30       <sup>1</sup>H NMR (free base, CDCl<sub>3</sub>) δ 1.10 (dd, 6H), 1.15-2.1 (m,  
6H), 2.65-2.90 (m+s, 5H), 3.05-3.25 (m+s, 4H), 3.35 (d, 1H),  
3.55 (d, 1H), 3.90 (d, 1H), 4.30 (m, 1H), 6.65 (d, 1H), 6.95  
(d, 1H), 7.05-7.4 (m, 6H).

EXAMPLE 16Cis-3-[2-methoxy-5-(N-isopropyl-N-methanesulfonyl-amino)benzylamino]-2-phenylpiperidine dihydrochloride

M.P. 268-270°C, 65% yield.



-54-

Anal. calc'd for  $C_{23}H_{33}N_3O_3S \cdot 2HCl$ : C, 54.75; H, 6.99; N, 8.32. Found: C, 54.75, H, 6.99, N, 8.29.

$^1H$  NMR (free base,  $CDCl_3$ )  $\delta$  1.10 (dd, 6H), 1.45 (d, 1H), 1.60 (tt, 1H), 1.7-1.95 (m, 3H), 2.12 (d, 1H), 2.80 (m, 2H), 2.90 (s, 3H), 3.25 (d, 1H), 3.35 (d, 1H), 3.50 (s, 3H), 3.70 (d, 1H), 3.90 (d, 1H), 4.50 (m, 1H), 6.65 (d, 1H), 6.90 (d, 1H), 7.05 (dd, 1H), 7.30 (m, 5H).

#### EXAMPLE 17

Cis-3-[2-methoxy-5-(N-methyl-N-trifluoromethanesulfonylamino)benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 245-250°C, 24% yield.

Anal. calc'd for  $C_{21}H_{26}F_3N_3O_3S \cdot 2HCl$ : C, 47.55; H, 5.32; N, 7.92. Found: C, 47.55, H, 5.32, N, 7.86.

$^1H$  NMR (free base,  $CDCl_3$ )  $\delta$  1.50 (d, 1H), 1.60 (tt, 1H), 1.8-2.00 (m, 3H), 2.15 (d, 1H), 2.85 (m, 2H), 3.25 (d, 1H), 3.35 (s, 3H), 3.40 (d, 1H), 3.50 (s, 3H), 3.65 (d, 1H), 3.90 (d, 1H), 6.65 (d, 1H), 6.98 (d, 1H), 7.10 (dd, 1H), 7.25 (m, 5H).

#### EXAMPLE 18

Cis-3-[2-methoxy-5-(N-thiazolidine-S,S-dioxide)-benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 263-265°C, 36% yield.

Anal. calc'd for  $C_{22}H_{29}N_3O_3S_2 \cdot 2HCl$ : C, 54.09; H, 6.40; N, 8.60. Found: C, 53.87, H, 6.43, N, 8.45.

$^1H$  NMR (free base,  $CDCl_3$ )  $\delta$  1.40 (d, 1H), 1.60 (tt, 1H), 1.75 (m, 2H), 1.90 (m, 1H), 2.15 (d, 1H), 2.50 (m, 2H), 2.80 (m, 2H), 3.2-3.50 (m, 7H), 3.55-3.70 (m, 3H), 3.90 (d, 1H), 6.65 (d, 1H), 6.95 (d, 1H), 7.1-7.40 (m, 6H).

#### EXAMPLE 19

Cis-3-[2-trifluoromethoxy-5-(N,N-bis(methanesulfonyl)amino)benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 256-257°C, 29% yield.

Anal. calc'd for  $C_{21}H_{26}F_3N_5O_6S_2 \cdot 2HCl$ : C, 42.43; H, 4.75; N, 7.07. Found: C, 42.38, H, 4.77, N, 6.94.

-55-

<sup>1</sup>H NMR (free base, CDCl<sub>3</sub>) δ 1.50 (d, 1H), 1.6-1.90 (m, 4H), 2.10 (d, 1H), 2.75-2.95 (m, 2H), 3.2-3.40 (m+s, 7H), 3.50 (d, 1H), 3.65 (d, 1H), 3.95 (d, 1H), 7.15-7.45 (m, 8H).

EXAMPLE 20

5     Cis-3-[2-methoxy-5-(N,N-diethylaminosulfonyl)-benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 267-269°C, 29% yield.

Anal. calc'd for C<sub>23</sub>H<sub>33</sub>N<sub>3</sub>O<sub>3</sub>S<sub>2</sub>•2HCl: C, 54.75; H, 6.99; N, 8.32. Found: C, 54.98; H, 7.34; N, 8.18.

10     <sup>1</sup>H NMR (free base, CDCl<sub>3</sub>) δ 1.15 (t, 6H), 1.50 (d, 1H), 1.6-2.00 (m, 4H), 2.10 (d, 1H), 2.80 (m, 2H), 3.15 (q, 4H), 3.30 (d, 1H), 3.55 (s+d, 4H), 3.70 (d, 1H), 3.95 (d, 1H), 6.70 (d, 1H), 7.30 (m, 5H), 7.65 (dd, 1H).

EXAMPLE 21

15     Cis-3-[2-trifluoromethoxy-5-(N-methyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 247-248°C, 43% yield.

20     Anal. calc'd for C<sub>21</sub>H<sub>26</sub>F<sub>3</sub>N<sub>3</sub>O<sub>3</sub>S<sub>2</sub>•2HCl: C, 47.55; H, 5.32; N, 7.92. Found: C, 47.51, H, 5.47, N, 7.60.

<sup>1</sup>H NMR (free base, CDCl<sub>3</sub>) δ 1.50 (d, 1H), 1.6-1.95 (m, 4H), 2.10 (d, 1H), 2.75 (s, 3H), 2.85 (m, 2H), 3.15 (s, 3H), 3.30 (d, 1H), 3.50 (d, 1H), 3.65 (d, 1H), 3.95 (d, 1H), 7.1-7.45 (m, 8H).

25

EXAMPLE 22

Cis-3-[2-isopropoxy-5-(N-methyl-N-trifluoromethanesulfonylamino)benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 267-273°C, 7% yield.

30     Anal. calc'd for C<sub>23</sub>H<sub>30</sub>F<sub>3</sub>N<sub>3</sub>O<sub>3</sub>S<sub>2</sub>•2HCl: C, 49.46; H, 5.41; N, 7.52. Found: C, 49.71, H, 5.72, N, 7.30.

<sup>1</sup>H NMR (free base, CDCl<sub>3</sub>) δ 1.15 (dd, 6H), 1.4-1.95 (m, 5H), 2.15 (d, 1H), 2.30 (m, 2H), 3.15-3.4 (m+s, 5H), 3.55 (d, 1H), 3.90 (d, 1H), 4.35 (m, 1H), 6.65 (d, 1H), 6.95 (d, 35 1H), 7.10 (dd, 1H), 7.30 (m, 5H).

-56-

EXAMPLE 23

Cis-3-[2-methoxy-5-(N-methyl-N-isopropylsulfonyl-amino)benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 264-266°C, 22% yield.

5      Anal. calc'd for  $C_{22}H_{33}N_3O_3S \cdot 2HCl$ : C, 54.75; H, 6.99; N, 8.32. Found: C, 54.91, H, 7.04, N, 8.23.

$^1H$  NMR (free base,  $CDCl_3$ ,  $\delta$ ) 1.35 (d, 6H), 1.45 (d, 1H), 1.55-1.95 (m, 4H), 2.15 (d, 1H), 2.85 (m, 2H), 3.25 (m+s, 5H), 3.35 (d, 1H), 3.50 (s, 3H), 3.65 (d, 1H), 3.90 (d, 1H),  
10 6.65 (d, 1H), 7.05 (d, 1H), 7.15-7.35 (m, 6H).

EXAMPLE 24

Cis-3-[2-cyclopentyloxy-5-(N-methyl-N-methanesulfonyl-amino)benzyl]amino-2-phenylpiperidine dihydrochloride hemihydrate

15      M.P. 252-255°C, 37% yield.

Anal. calc'd for  $C_{25}H_{33}N_3O_3S \cdot 2HCl \cdot 1/2H_2O$ : C, 55.65, H, 7.10, N, 7.79. Found: C, 55.51, H, 6.95, N, 7.73.

$^1H$  NMR (free base,  $CDCl_3$ )  $\delta$  1.4-1.95 (m, 13H), 2.10 (d, 1H), 2.7-2.90 (m+s, 5H), 3.20 (s, 3), 3.25 (d, 1H), 3.35 (d, 1H), 3.55 (d, 1H), 3.85 (d, 1H), 4.55 (m, 1H), 6.65 (d, 1H),  
20 6.95 (d, 1H), 7.10 (dd, 1H), 7.25 (m, 5H).

EXAMPLE 25

Cis-3-[2-methoxy-5-(N-methyl-N-(4-methylphenylsulfonyl)-amino)benzyl]amino-2-phenylpiperidine dihydrochloride

25      M.P. 215-220°C, 42% yield.

Anal. calc'd for  $C_{27}H_{33}N_3O_3S \cdot 2HCl$ : C, 58.69, H, 6.38, N, 7.60. Found: C, 58.46, H, 6.30, N, 7.41.

$^1H$  NMR (free base,  $CDCl_3$ ,  $\delta$ ) 1.30-2.04 (m, 7H), 2.40 (s, 3H), 2.74 (m, 2H), 3.05 (s, 3H), 3.25 (d, 1H), 3.40 (s, 3H),  
30 3.52 (d, 1H), 3.80 (d, 1H), 6.52 (d, 1H), 6.62 (d, 1H), 6.85 (dd, 1H), 7.10-7.42 (m, 9H).

EXAMPLE 26

Cis-3-[2-isopropoxy-5-(N-methyl-N-(4-methylphenylsulfonyl)amino)benzyl]amino-2-phenylpiperidine dihydrochloride

35

M.P. 215-219°C, 3.2% yield.

Anal. calc'd for  $C_{29}H_{37}N_3O_3S \cdot 2HCl$ : C, 59.99, H, 6.77, N,

-57-

7.23. Found: C, 59.98, H, 6.83, N, 7.19.

<sup>1</sup>H NMR (free base, CDCl<sub>3</sub>, δ) 1.04 (dd, 6H), 1.30-2.05 (m, 7H), 2.40 (s, 3H), 2.75 (m, 2H), 3.04 (s, 3H), 3.24 (d, 1H), 3.44 (d, 1H), 3.80 (d, 1H), 4.26 (m, 1H), 6.55 (d, 1H),  
5 6.63 (d, 1H), 6.85 (dd, 1H), 7.10-7.42 (m, 9H).

EXAMPLE 27

Cis-3-[2-isopropoxy-5-(N-isopropyl-N-methanesulfonyl-amino)benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 243-246°C, 23% yield.

10 Anal. calc'd for C<sub>25</sub>H<sub>37</sub>N<sub>3</sub>O<sub>3</sub>S•2HCl: C, 56.38, H, 7.38, N, 7.89. Found: C, 56.52, H, 7.03, N, 7.70.

<sup>1</sup>H NMR (free base, CDCl<sub>3</sub>, δ) 1.10-1.5 (dd+dd, 12H), 1.40-2.20 (m, 6H), 2.60 (m, 2H), 2.80 (s, 3H), 3.30 (m, 1H), 3.35 (d, 1H), 3.65 (d, 1H), 3.80 (d, 1H), 4.35 (m, 1H), 4.50  
15 (m, 1H), 6.95 (d, 1H), 7.05 (dd, 1H), 7.30 (m, 5H).

EXAMPLE 28

Cis-3-[2-isopropoxy-5-(N,N-diethylaminosulfonyl)-benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 246-248°C (dec.), 98% yield.

20 Anal. calc'd for C<sub>25</sub>H<sub>37</sub>N<sub>3</sub>O<sub>3</sub>S•2HCl: C, 56.39, H, 7.38, N, 7.89. Found: C, 56.29, H, 7.29, N, 7.82.

<sup>1</sup>H NMR (free base, CDCl<sub>3</sub>, δ) 1.11 (m, 12H), 1.37-2.15 (m, 6H), 2.72-2.83 (m, 2H), 3.12-3.28 (q+m, 5H), 3.33 (d, 1H), 3.60 (d, 1H), 3.85 (d, J=2.2 Hz, 1H), 4.38 (m, 1H),  
25 6.71 (d, 1H), 7.25 (m, 5H), 7.48 (d, 1H), 7.57 (dd, 1H).

EXAMPLE 29

Cis-3-[2-methoxy-5-(N-methyl-N-phenylmethylsulfonyl-amino)benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 266-269°C (dec.), 23% yield.

30 Anal. calc'd for C<sub>27</sub>H<sub>33</sub>N<sub>3</sub>O<sub>3</sub>S•2HCl: C, 58.69, H, 6.39, N, 7.60. Found: C, 58.70, H, 6.54, N, 7.41.

<sup>1</sup>H NMR (free base, CDCl<sub>3</sub>, δ) 1.40-2.30 (m, 6H), 2.80 (m, 2H), 3.07 (s, 3H), 3.30 (m, 1H), 3.35 (d, 1H), 3.50 (s, 3H), 3.65 (d, 1H), 3.90 (d, 1H), 4.20 (s, 2H), 6.62 (d, 1H), 6.90  
35 (d, 1H), 7.08 (dd, 1H), 7.20-7.45 (m, 10H).

-58-

EXAMPLE 30

Cis-3-[(2,3-dihydro-5-methoxy-1-methanesulfonyl-6-indolyl)methylamino]-2-phenylpiperidine dihydrochloride

M.P. 255-258°C, 27% yield.

5     Anal. calc'd for  $C_{22}H_{29}N_3O_3S \cdot 2HCl$ : C, 54.09, H, 6.40, N, 8.60. Found: C 54.10, H, 6.21, N, 8.52.

$^1H$  NMR (free base,  $CDCl_3$ ,  $\delta$ ) 1.35-2.20 (m, 7H), 2.75 (m, 1H), 2.80 (s, 3H), 3.05 (t, 2H), 3.25 (m, 1H), 3.35 (d, 1H), 3.40 (s, 3H), 3.60 (d, 1H), 3.95 (m, 3H), 6.55 (s, 1H), 7.15  
10 (s, 1H), 7.30 (m, 5H).

EXAMPLE 31

(1SR,2SR,3SR,4RS)-3-(2-methoxy-5-(N-methyl-N-methanesulfonylamino)benzyl)-amino-2-benzhydryl-[2.2.1]-azanorbornane dihydrochloride monohydrate

15     M.P. 196-200°C.

Anal. calc'd for  $C_{29}H_{35}N_3O_3S \cdot 2HCl \cdot H_2O$ : C, 58.38, H, 6.59; N, 7.04. Found: C, 58.71; H, 6.52; N, 6.93.

$^1H$  NMR ( $D_2O$ ,  $\delta$ ) 1.85 (m, 1H), 2.35 (m, 1H), 3.06 (s, 3H), 3.27-3.63 (m+s+s, 10H), 3.85 (d, 1H), 3.96 (d+d, 2H),  
20 4.26 (d, 1H), 4.39 (d, 1H), 4.8 (s,  $D_2O$ ), 5.16 (m, 1H), 6.97 (d, 1H), 7.21 (d, 1H), 7.31-7.50 (m, 1H).

EXAMPLE 32

(1SR,2SR,3SR,4RS)-3-(2-isopropoxy-5-(N-methyl-N-methanesulfonylamino)benzyl)amino-2-benzhydryl-[2.2.1]-  
25 azanorbornane dihydrochloride monohydrate

M.P. 182-183°C.

Anal. calc'd for  $C_{31}H_{39}N_3O_3S \cdot 2HCl \cdot H_2O$ : C, 53.54; H, 5.58; N, 6.46. Found: C, 53.36; H, 5.71; N, 6.40.

$^1H$  NMR ( $D_2O$ ,  $\delta$ ) 1.20 (t, 6H), 1.90 (m, 1H), 2.35 (m,  
30 1H), 3.06 (s, 3H), 3.26 (s, 3H), 3.29-3.47 (m, 4H), 3.84 (m, 3H), 4.14 (d, 1H), 4.36 (d, 1H), 4.45 (m, 1H), 4.80 (s,  $D_2O$ ), 5.08 (m, 1H), 6.96-7.04 (m, 2H), 7.26-7.47 (m, 11H).

EXAMPLE 33

(1SR,2SR,3SR,4RS)-3-(2-methoxy-5-(N-methyl-N-trifluoromethanesulfonylamino)benzyl)amino-2-benzhydryl-[2.2.1]-  
35 azanorbornane dihydrochloride

M.P. 186°C.

-59-

HRMS calc'd for  $C_{29}H_{32}F_3N_3O_3S$ : 559.2116. Found:  
559.2197.

$^1H$  NMR ( $D_2O$ ,  $\delta$ ) 1.85 (m, 1H), 2.34 (m, 1H), 3.36-3.55  
(m+s, 10H), 3.72-3.85 (d+d, 4H), 4.14 (d, 1H), 4.37 (d, 1H),  
5 4.80 (s,  $D_2O$ ), 5.03 (m, 1H), 6.97 (d, 1H), 7.24 (d, 1H),  
7.32-7.53 (m, 11H).

#### EXAMPLE 34

(1SR,2SR,3SR,4RS)-3-(2-methoxy-5-(N-methyl-N-phenylmethanesulfonylamino)benzyl)amino-2-benzhydryl-  
10 [2.2.1]-azanorbornane dihydrochloride hydrate

M.P. 178°C.

Anal. calc'd for  $C_{35}H_{39}N_3O_3S \cdot 2HCl \cdot 1.5H_2O$ : C, 58.76; H,  
7.00; N, 6.63. Found: C, 59.15; H, 6.60; N, 6.40.

$^1H$  NMR ( $D_2O$ ,  $\delta$ ) 1.81 (m, 1H), 2.32 (m, 1H), 3.24-3.37  
15 (m, 8H), 3.51 (m, 3H), 3.68 (m, 2H), 3.79 (d, 1H), 3.95 (d,  
1H), 4.35 (d, 1H), 4.62 (s, 1H), 4.82 (s+m, 1H), 4.97 (m,  
1H), 6.69 (d, 1H), 6.85 (d, 1H), 7.11 (dd, 1H), 7.37-7.50  
(m, 15).

#### EXAMPLE 35

20 (1SR,2SR,3SR,4RS)-3-(2-methoxy-5-(N-isopropyl-N-methanesulfonylamino)benzyl)amino-2-benzhydryl-[2.2.1]-  
azanorbornane dihydrochloride

M.P. 238°C (dec.).

Anal. calc'd for  $C_{31}H_{39}N_3O_3S \cdot 2HCl$ : C, 61.08; H, 6.49; N,  
25 6.74; N, 6.74. Found: C, 61.38; H, 6.81; N, 6.93.

$^1H$  NMR ( $D_2O$ ,  $\delta$ ) 1.14 (d, 6H), 1.87 (m, 1H), 2.38 (m,  
1H), 3.18 (s, 3H), 3.34-3.61 (m+s, 7H), 3.89 (d, 1H), 4.05  
(m, 2H), 4.31-4.46 (m, 3H), 4.8 (s,  $D_2O$ ), 5.19 (m, 1H), 7.01  
(d, 1H), 7.20 (d, 1H), 7.34-7.52 (m, 11H).

30

#### EXAMPLE 36

(1SR,2SR,3SR,4RS)-3-(2-methoxy-5-(1,1-dioxo-2-isothiazolidinyl)benzyl)amino-2-benzhydryl-[2.2.1]-  
azanorbornane dihydrochloride

M.P. 206-207°C.

35 Anal. calc'd for  $C_{30}H_{35}N_3O_3S \cdot 2HCl$ : C, 60.09; H, 6.39; N,  
7.01. Found: C, 59.77; H, 6.15; N, 6.94.

-60-

<sup>1</sup>H NMR (D<sub>2</sub>O, δ) 1.90 (m, 1H), 2.35 (m, 1H), 2.56 (m, 2H), 3.33-3.62 (m+s, 10H), 3.77-3.83 (m, 4H), 3.96 (d, 1H), 4.15 (d, 1H), 4.41 (d, 1H), 4.8 (s, D<sub>2</sub>O), 5.10 (m, 1H), 7.00 (d, 1H), 7.13 (d, 1H), 7.32-7.47 (m, 11H).

5

EXAMPLE 37

(1SR,2SR,3SR,4RS)-3-[(2,3-dihydro-5-methoxy-1-methanesulfonyl-6-indolyl)methylamino]benzyl]-2-benzhydryl-[2.2.1]-azanorbornane dihydrochloride

M.P. 250°C.

10

Anal. calc'd for C<sub>30</sub>H<sub>35</sub>N<sub>3</sub>O<sub>3</sub>S•2HCl: C, 63.34; H, 6.38; N, 6.33. Found: C, 63.48; H, 6.15; N, 6.32.

<sup>1</sup>H NMR (D<sub>2</sub>O, δ) 1.90 (m, 1H), 2.38 (m, 1H), 2.99 (s, 3H), 3.20 (t, 2H), 3.33-3.55 (m+s, 8H), 3.86 (d, 1H), 3.97-4.06 (m, 4H), 4.19 (d, 1H), 4.39 (d, 1H), 4.82 (s, D<sub>2</sub>O), 5.13 (m, 1H), 6.96 (s, 1H), 7.12 (s, 1H), 7.36-7.51 (m, 10H).

15

EXAMPLE 38

(2S,3S)-N-(2-Methoxy-5-methylthiophenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine Mesylate

A. (2S,3S)-2-Diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine

20

(2S,3S)-N-(2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine (4.13 g, 10 mmol) was hydrogenated in methanol (MeOH) (40 ml)/6N HCl (10 ml) by using 20% palladium hydroxide on carbon (0.2 g) at 2.5 kg/cm<sup>2</sup> of hydrogen for 60 hours. The filtrate was concentrated and the residue was partitioned between 2N NaOH and CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was dried over MgSO<sub>4</sub>, and concentrated to give the crude product, which was recrystallized from ethanol (EtOH) to afford the pure title compound (2.80 g, 96%).

25

B. 2-Methoxy-5-methylthiobenzaldehyde

30

2-(2-Methoxy-5-methylthiophenyl)-1,3-dioxolane (2.40 g, 10 mmol) was stirred in 1N HCl (2 ml)/acetone (30 ml). After the starting material disappeared (ca. 2 hours), the solution was concentrated. The residue was partitioned between methylene chloride (CH<sub>2</sub>Cl<sub>2</sub>) and saturated sodium bicarbonate (NaHCO<sub>3</sub>) solution. The organic layer was washed

35

-61-

with H<sub>2</sub>O, dried over MgSO<sub>4</sub>, and evaporated to give the aldehyde. (1.96 g, 100%).

C. (2S,3S)-N-(2-Methoxy-5-methylthiophenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine Mesylate

5 To a solution of a 2-methoxy-5-methylthiobenzaldehyde (765 mg, 4.2 mmol) and (2S,3S)-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine (1170 mg, 4 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (40 ml) was added in portions sodium triacetoxyborohydride (933 mg, 4.4 mmol). The mixture was stirred until the amine  
10 disappeared. The solution was carefully neutralized with an ice cooled saturated NaHCO<sub>3</sub> solution. The organic layer was washed with H<sub>2</sub>O, dried over MgSO<sub>4</sub>, and concentrated to give the product (1.61 g, 88%). To the solution of the product in acetone was added one equivalent methanesulfonic acid.  
15 Then the precipitated mesylate salt was collected (1.51 g, 66%).

M.P. 234°C.

IR (KBr) cm<sup>-1</sup>: 3400, 2950, 1630, 1600, 1490, 1455, 1240, 1210, 1195, 1060, 785, 750, 710.

20 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.40 (1H, br), 7.5-7.2 (10 H, m), 7.17 (1H, d, J=8.4 Hz), 6.69 (1H, d, J=8.4 Hz), 6.66 (1H, br, s), 4.56 (1H, d, J=12.1 Hz), 4.25 (1H, m), 3.70-3.35 (5H, m), 3.55 (3H, s), 3.30-3.15 (2H, m), 2.46 (3H, s), 2.42 (3H, s), 2.25 (1H, m), 2.05 (1H, m), 2.00-1.60 (3H, m).

25 EXAMPLE 39

(2S,3S)-N-(2-Methoxy-5-methylsulfinylphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine Hydrochloride

A solution of (2S,3S)-N-(2-methoxy-5-methylthiophenyl)-  
30 methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine (180 mg, 0.392 mmol) in MeOH (20 ml) was added to a solution of sodium periodate (NaIO<sub>4</sub>) (92 mg, 0.432 mmol) in H<sub>2</sub>O (10 ml). The mixture was stirred for 24 hours. The precipitate (NaIO<sub>3</sub>) was filtered off. The filtrate was concentrated and  
35 the residue was partitioned between H<sub>2</sub>O and CH<sub>2</sub>Cl<sub>2</sub> (20 ml). The water layer was extracted twice with CH<sub>2</sub>Cl<sub>2</sub>. The combined CH<sub>2</sub>Cl<sub>2</sub> was dried over MgSO<sub>4</sub> and concentrated to give



-62-

the sulfoxide, which was converted to HCl salt by using HCl-ether. (Yield, 180 mg, 97%).

M.P. 183°C.

IR (KBr)  $\text{cm}^{-1}$ : 3420, 3190, 1605, 1495, 1455, 1260, 1020,  
5 755, 710.

$^1\text{H}$  NMR ( $\text{CDCl}_3$  + DMSO)  $\delta$ : 8.11 (1H, br), 8.00 (1H, br),  
7.70 (2H, m), 7.65 (1H, m), 7.44-7.20 (7H, m), 6.92 (1H, m),  
6.48 (1H, br), 5.49 (1H, m), 4.45 (1H, br), 4.20 (2H, m),  
3.95 (1H, m), 3.16 (1.5H, s), 3.12 (1.5H, s), 3.15 (2H, m),  
10 2.80 (1.5H, s), 2.77 (1.5H, s), 2.85-2.50 (5H, m), 2.15-1.85  
(2H, m).

#### EXAMPLE 40

(2S,3S)-N-(5-Ethylthio-2-methoxyphenyl)methyl-2-  
diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine  
15 Hydrochloride

The title compound was obtained using the same procedure as described in Example 38, except that 5-ethylthio-2-methoxybenzaldehyde was substituted for 2-methoxy-5-methylthiobenzaldehyde. The yield of the product  
20 was 76%.

M.P. 254°C.

IR (KBr)  $\text{cm}^{-1}$ : 3450, 3190, 2950, 1490, 1455, 1250, 1030,  
715.

$^1\text{H}$  NMR (DMSO)  $\delta$ : 7.97 (1H, br), 7.68 (2H, m), 7.51 (2H,  
25 m), 7.50-6.85 (9H, m), 5.46 (2H, m), 4.25-3.30 (4H, m), 3.44  
(3H, s), 3.16 (2H, m), 2.89 (2H, q, 7.3 Hz), 2.65 (1H, m),  
2.30 (1H, m), 2.15-1.80 (4H, m), 1.19 (3H, t, 7.3Hz).

#### EXAMPLE 41

(2S,3S)-N-(5-Trifluoroacetylamino-2-methoxyphenyl)-  
methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine  
30 Mesylate

The title compound was obtained using the same procedure as described in Example 38, except that 5-trifluoroacetylamino-2-methoxybenzaldehyde was substituted  
35 for 2-methoxy-5-methylthiobenzaldehyde. The yield of the product was 96%.

M.P. 148°C.

-63-

IR (KBr)  $\text{cm}^{-1}$ : 3430, 3050, 1610, 1500, 1200, 1060, 750, 710, 565.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 9.50 (1H, br), 7.80 (1H, m), 7.5-7.1 (12H, m), 6.68 (1H, d,  $J=9.2$  Hz), 4.68 (1H, m), 4.49 (1H, m), 3.80-3.50 (2H, m), 3.52 (3H, s), 3.50-3.20 (5H, m), 2.48 (3H, s), 2.42 (1H, m), 2.23 (1H, m), 1.99 (2H, m), 1.71 (1H, m).

#### EXAMPLE 42

##### 10 (2S,3S)-N-(2-Methoxy-5-dimethylaminophenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine Mesylate

The title compound was obtained using the same procedure as described in Example 38, except that 2-methoxy-5-dimethylaminobenzaldehyde was substituted for 2-methoxy-5-methylthiobenzaldehyde. The yield of the product was 75%.

15 M.P. 240°C.

IR (KBr)  $\text{cm}^{-1}$ : 3420, 2960, 1620, 1510, 1455, 1240, 1210, 1195, 1060, 785, 750, 710.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 8.37 (1H, br), 7.45-7.20 (10H, m), 6.67 (2H, m), 6.38 (1H, m), 4.60 (1H, m), 4.23 (1H, m), 20 3.30-3.70 (5H, m), 3.49 (3H, s), 3.10-3.35 (2H, m), 2.86 (6H, s), 2.51 (3H, s), 2.42 (1H, m), 2.26 (1H, m), 2.15-1.50 (3H, m).

#### EXAMPLE 43

##### 25 (2S,3S)-N-(5-Amino-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine Mesylate

(2S,3S)-N-(5-Trifluoroacetyl-amino-2-methoxyphenyl)-methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine (1.52 g, 3 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 ml)/saturated  $\text{NaHCO}_3$  (20 ml) was stirred vigorously for 8 hours. The  $\text{CH}_2\text{Cl}_2$  layer was 30 washed with water, dried over  $\text{MgSO}_4$ , and concentrated to give the title compound, which was converted to HCl salt by using HCl-ether. (Yield, 1.35 g, 81%).

M.P. 237°C.

IR (KBr)  $\text{cm}^{-1}$ : 3430, 2900, 1625, 1505, 1455, 1270, 1020, 35 755, 710.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : (free base) 7.45-7.05 (10H, m), 6.55 (1H, m), 6.47 (1H, m), 5.79 (1H, m), 4.50 (1H, d, 12 Hz),

-64-

3.70 (1H, m), 3.52 (3H, s), 3.50 (1H, d, 14 Hz), 3.28 (1H, d, 14 Hz), 3.20 (1H, m), 2.92 (1H, m), 2.79 (2H, m), 2.61 (1H, m), 2.04 (1H, m), 1.91 (1H, m), 1.65 (1H, m), 1.55 (1H, m), 1.28 (1H, m).

5

EXAMPLE 44(2S,3S)-N-(2-Methoxy-5-methylsulfonylphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine Mesylate

The title compound of Example 38 (free amine) (1.20 g, 2.62 mmol) was treated with methanolic HCl to give the  
10 hydrochloride salt. Evaporation of the solvent gave crude (2S,3S)-N-(2-methoxy-5-methylthiophenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine dihydrochloride.

To a stirred and ice-cooled solution of (2S,3S)-N-(2-  
15 methoxy-5-methylthiophenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine dihydrochloride in methanol (25 mL) was added a solution of oxone (2.41 g) in water (25 mL). The reaction mixture was stirred at room temperature for 2.5 hours. The reaction mixture was basified to pH 10-  
20 11 with 1N NaOH aq. solution with ice-cooling, and extracted with CHCl<sub>3</sub> (80 mL X 4). The combined organic layers were washed with brine (80 mL), dried (MgSO<sub>4</sub>) and concentrated in vacuo to give crude (2S,3S)-N-(2-methoxy-5-methylsulfonylphenyl)methyl-2-diphenylmethyl-1-  
25 azabicyclo[2.2.2]octan-3-amine (white soap, 1.49 g). The residue was purified by chromatography on silica gel (60 g) with chloroform-methanol (20:1-10:1) to give (2S,3S)-N-(2-methoxy-5-methylsulfonylphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine (1.08 g, 79%) as a white  
30 amorphous solid.

To a solution of (2S,3S)-N-(2-methoxy-5-methylsulfonylphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine (400 mg, 0.82 mmol) in acetone (10 mL) was added methanesulfonic acid (0.41 mmol, 39.2 mg). The precipitated white solid was filtered off to  
35 give the title compound (218 mg, 30.3%, 1st crop).

M.P. 240-241°C.

-65-

IR (KBr, free amine): 3430, 2940, 1597, 1493, 1449, 1350, 1306, 1256, 1186, 1128, 960, 820, 800, 754, 704  $\text{cm}^{-1}$ .

$^1\text{H}$  NMR (270 MHz,  $\text{CDCl}_3$ , ppm) (free amine): 7.77 (1H, dd,  $J=2.6, 8.4$  Hz), 7.50 (1H, d,  $J=2.6$  Hz), 7.37-7.03 (10H, m),  
5 6.81 (1H, d,  $J=8.4$  Hz), 4.47 (1H, d,  $J=12.1$  Hz), 3.71 (1H, dd,  $J=7.7, 12.1$  Hz), 3.62 (3H, s), 3.61 (1H, d,  $J=13.6$  Hz),  
3.21 (1H, d,  $J=13.6$  Hz), 3.28-3.10 (1H, m), 3.01 (3H, s),  
2.94 (1H, dd,  $J=4.4, 7.7$  Hz), 2.83-2.74 (2H, m), 2.63 (1H, br.t,  $J=11.7$  Hz), 2.10-2.03 (1H, m), 1.95-1.45 (3H, m),  
10 1.35-1.20 (1H, m).

Anal. calc'd for  $\text{C}_{29}\text{H}_{34}\text{N}_2\text{O}_2\text{S}\cdot\text{CH}_3\text{SO}_3\text{H}\cdot 2\text{H}_2\text{O}$ : C, 57.86; H, 6.80%; N, 4.50%. Found: C, 57.93%; H, 6.97%; N, 4.34%.

#### EXAMPLE 45

#### cis-2-(Diphenylmethyl)-N-((5-amino-2-methoxyphenyl)- 15 methyl)-1-azabicyclo[2.2.2]octan-3-amine

To a 50 mL round-bottomed flask equipped with Dean-Stark trap, condenser and  $\text{N}_2$  inlet were added 430 mg (2.38 mmol) 2-methoxy, 5-nitrobenzaldehyde, 578 mg<sup>-</sup> (1.98 mmol) cis-2-(diphenylmethyl)-1-azabicyclo[2.2.2]octan-3-amine,  
20 4 mg camphorsulfonic acid, and 10 mL toluene. The reaction was refluxed with removal of water for 14 hours, then cooled and evaporated. The residue was dissolved in 10 mL tetrahydrofuran, treated with 5 ml (10 mmol) of a 2.0 M solution of borane/methyl sulfide in tetrahydrofuran, and  
25 refluxed for 3 days. The reaction was then cooled and evaporated, taken up to 10 mL ethanol, treated with 1 g sodium carbonate and 1 g cesium fluoride, and refluxed for 2 days. The reaction was cooled, partitioned between water and methylene chloride, and the organic layer was separated,  
30 washed with brine, dried over sodium sulfate, and evaporated. The residue was chromatographed on silica gel using acetonitrile/water/acetic acid as eluant, and the product fractions were isolated to afford 347 mg (41%) of an amorphous solid, which crystallized from isopropanol to give  
35 M.P. 164-169°C.

$^1\text{H}$  NMR ( $\delta$ ,  $\text{CDCl}_3$ ): 1.23 (m, 1H), 1.49 (m, 1H), 1.60 (m, 1H), 1.90 (m, 1H), 2.03 (m, 1H), 2.60 (m, 2H), 2.75 (m, 2H),

-66-

2.89 (m, 1H), 3.20 (m, 1H), 3.39 (ABq,  $J_{AB}=16$ ,  $\Delta\nu=62$ , 2H), 3.51 (s, 3H), 3.66 (dd,  $J=8,12$ , 1H), 4.49 (d,  $J=12$ , 1H), 5.78 (m, 1H), 6.4-6.6 and 7.0-7.4 (m, 13H).

$^{13}\text{C}$  NMR ( $\delta$ ,  $\text{CDCl}_3$ ): 20.1, 24.8, 25.6, 42.1, 45.9, 49.3, 53.7, 54.3, 56.0, 61.8, 111.5, 114.0, 116.6, 125.9, 126.3, 127.6, 128.4, 129.0, 129.1, 139.7, 143.6, 145.7, 150.6.

IR ( $\text{cm}^{-1}$ , KBr): 1620 and 1580.

MS (%): 428 (parent + 1, 1), 291 (22), 260 (100), 136 (54), 106 (23).

Anal. calc'd for  $\text{C}_{23}\text{H}_{33}\text{N}_3\text{O}$ : C 78.65, H 7.78, N 9.83.  
Found: C 78.73, H 7.87, N 9.71.

The title compounds of Example 46 to 58 were prepared by a procedure similar to that described in Example 9.

#### EXAMPLE 46

(5-Isopropylsulfonyl-2-methoxybenzyl)-(2-phenylpiperidin-3-yl)amine dihydrochloride

17% yield, m.p. 278-280°C (dec.).

MS: m/e 402 ( $\text{M}^+$ ), 398, 283, 275.

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , free base)  $\delta$  1.25 (dd, 6H), 1.35-2.2 (m, 6H), 2.8 (m, 2H), 3.15 (m, 1H), 3.25 (d, 1), 3.35 (d, 1H), 3.5 (s, 3H), 3.65 (d, 1H), 3.9 (d, 1H), 6.75 (d, 1H), 7.25 (m, 5H), 7.55 (s, 1H), 7.65 (dd, 1H).

Anal. calc'd for  $\text{C}_{22}\text{H}_{30}\text{N}_2\text{O}_3\text{S}\cdot 2\text{HCl}$ : C, 55.57; H, 6.78; N, 5.89. Found: C, 55.24; H, 6.54; N, 5.87.

#### EXAMPLE 47

N-Cyclopentyl-N-[4-methoxy-3-(2-phenylpiperidin-3-ylaminomethyl)phenyl]methanesulfonamide dihydrochloride hemihydrate

30% yield, m.p. 249-252°C.

FABMS: m/e 458 ( $\text{M}^{+1}$ , 100%), 282 (10), 160 (55%).

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , free base)  $\delta$  1.25-1.65 (m, 8H), 1.75-2.05 (m, 5H), 2.15 (d, 1H), 2.8 (m, 2H), 2.9 (s, 3H), 3.25 (d, 1H), 3.35 (d, 1H), 3.5 (s, 3H), 3.7 (d, 1H), 3.9 (d, 1H), 4.45 (m, 1H), 6.65 (d, 1H), 6.9 (d, 1H), 7.05 (dd, 1H), 7.25 (m, 5H).

Anal. calc'd for  $\text{C}_{25}\text{H}_{35}\text{N}_3\text{O}_3\text{S}\cdot 2\text{HCl}\cdot 1/2\text{H}_2\text{O}$ : C, 55.65; H, 7.10; N, 7.79. Found: C, 55.69; H, 6.55; N, 7.78.

-67-

EXAMPLE 48

N-Cyclohexylmethyl-N-(4-methoxy-3-(2-phenylpiperidin-3-ylaminomethyl)phenylmethanesulfonamide dihydrochloride hydrate

5 21% yield, m.p. 255-256°C (dec.).

FABMS: m/e 486 ( $M^{+1}$ ), 408.

$^1H$  NMR ( $CDCl_3$ , free base)  $\delta$  0.9-2.2 (m, 17H), 2.7-2.9 (m, 5H), 3.2-3.5 (m, 5H), 3.5 (s, 3H), 3.6 (d, 1H), 3.7 (d, 1H), 3.9 (d, 1H), 6.7 (d, 1H), 7.0 (d, 1H), 7.3 (dd, 1H),  
10 7.4 (m, 5H).

Anal. calc'd for  $C_{27}H_{39}N_3O_3S \cdot 2HCl \cdot 3/4H_2O$ : C, 56.68; H, 7.49; N, 7.34. Found: C, 56.63; H, 7.11; N, 7.59.

EXAMPLE 49

(5-Methoxy-2-methyl-1-methylsulfonyl-2,3-dihydro-1H-indol-6-ylmethyl)-(2-phenylpiperidin-3-yl)amine dihydrochloride

16% yield, m.p. 257-259°C.

FABMS: m/e 430 ( $M^{+1}$ , 10%), 254 (100%).

$^1H$  NMR ( $CDCl_3$ , free base)  $\delta$  1.45 (dd, 3H), 1.65 (t, 1H),  
20 1.8-2.2 (m, 4H), 2.6 (m, 1H), 2.75 (d, 3H), 2.85 (m, 1H), 3.3 (m, 1H), 3.4 (d, 3H), 3.45 (m, 1H), 3.65 (m, 1H), 3.9 (d, 1H), 4.4 (m, 1H), 6.55 (d, 1H), 7.15 (d, 1H), 7.25 (m, 5H).

Anal. calc'd for  $C_{23}H_{31}N_3O_3S \cdot 2HCl$ : C, 54.97; H, 6.22; N, 8.36. Found: C, 54.76; H, 6.45; N, 8.20.

EXAMPLE 50

1-[5-Methoxy-6-(2-phenylpiperidin-3-ylaminomethyl)-2,3-dihydroindol-1-yl]-heptan-1-one dihydrochloride hemihydrate

7% yield, m.p. 170-172°C.

30 FABMS: m/e 450 ( $M^{+1}$ , 100%), 274, 160.

$^1H$  NMR ( $CDCl_3$ , free base)  $\delta$  0.9 (t, 3H), 1.25-1.45 (m, 6H), 1.5-1.8 (m, 3H), 1.85-2.25 (m, 4H), 2.4 (t, 2H), 2.8 (m, 2H), 3.15 (t, 2H), 3.25 (m, 1H), 3.3 (s, 3H), 3.35 (d, 1H), 3.7 (d, 1H), 3.9 (d, 1H), 4.05 (t, 2H), 6.5 (s, 1H),  
35 7.25 (m, 5H), 8.0 (s, 1H).

Anal. calc'd for  $C_{28}H_{39}N_3O_2 \cdot 2HCl \cdot 1/2H_2O$ : C, 63.27; H, 7.96; N, 7.90. Found: C, 63.33; H, 8.51; N, 8.19.

-68-

EXAMPLE 51

2,4-Dimethylthiazole-5-sulfonic acid [4-methoxy-3-(2-phenylpiperidin-3-ylaminomethyl)phenyl]-methanamide dihydrochloride hemihydrate

5        24% yield, m.p. 260-264°C.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, free base) δ 1.3-2.0 (m, 5H), 2.05 (d, 1H), 2.15 (s, 3H), 2.7 (s, 3H), 2.8 (m, 2H), 3.15 (s, 3H), 3.25 (d, 1H), 3.35 (d, 1H), 3.45 (s, 3H), 3.6 (d, 1H), 3.85 (d, 1H), 6.55 (d, 1H), 6.8 (d, 1H), 6.95 (dd, 1H), 7.25 (m, 5H).

Anal. calc'd for C<sub>25</sub>H<sub>32</sub>N<sub>4</sub>O<sub>3</sub>•2HCl•1/2H<sub>2</sub>O: C, 51.54; H, 6.06; N, 9.62. Found: C, 51.31; H, 5.79; N, 9.76.

EXAMPLE 52

N-(4,5-dimethylthiazol-2-yl)-N-[4-methoxy-3(2-phenylpiperidin-3-ylaminomethyl)phenyl]methanesulfonamide dihydrochloride hemihydrate

40% yield, m.p. 247-249°C.

FABMS: m/e 501 (M<sup>+</sup>), 421, 381, 247 (100%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, free base) δ 1.4 (d, 1H), 1.6 (t, 1H), 1.75 (m, 2H), 1.9 (m, 1H), 2.15 (d, 1H), 2.3 (m, 6H), 2.85 (m, 2H), 3.25 (d, 1H), 3.35 (d+s, 4H), 3.55 (s, 3H), 3.7 (d, 1H), 3.9 (d, 1H), 6.7 (d, 1H), 7.15 (d, 1H), 7.25 (m, 6H).

Anal. calc'd for C<sub>25</sub>H<sub>32</sub>N<sub>4</sub>O<sub>3</sub>S<sub>2</sub>•2HCl•1/2H<sub>2</sub>O: C, 51.54; H, 6.06; N, 9.62. Found: C, 51.87; H, 5.81; N, 9.55.

25        EXAMPLE 53

{5-[(4,5-dimethylthiazol-2-yl)methylamino]-2-methoxybenzyl}-(2-phenylpiperidin-3-yl)amine trihydrochloride hydrate

26% yield, m.p. 220-225°C.

30        MS: m/e 436 (M<sup>+</sup>, 16%), 317 (45%), 262 (100%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, free base) δ 1.5 (m, 1H), 1.6 (m, 1H), 1.9 (m, 1H), 2.1 (s, 3H), 2.2 (s, 3H), 2.8 (m, 2H), 3.2 (m, 1H), 3.3 (s, 3H), 3.4 (d, 1H), 3.5 (s, 3H), 3.6 (d, 1H), 3.9 (d, 1H), 6.4 (d, 1H), 6.9 (d, 1H), 7.1 (q, 1H), 7.4 (m, 5H).

35        Anal. calc'd for C<sub>25</sub>H<sub>32</sub>N<sub>4</sub>OS•HCl•3/2H<sub>2</sub>O: C, 52.40; H, 6.68; N, 9.78. Found: C, 52.12; H, 6.64; N, 9.55.

-69-

EXAMPLE 54

{5-[(4,5-dimethylthiazol-2-ylamino)-2-methoxybenzyl]-  
(2-phenylpiperidin-3-yl)amine trihydrochloride

28% yield, m.p. 272-275°C.

5 MS: m/e 422 (M<sup>+</sup>, 40%), 303 (54%), 248 (100%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, free base) δ 1.35-2.15 (m, 7H), 2.18 (s, 3H), 2.23 (s, 3H), 2.8 (m, 2H), 3.28 (d, 1H), 3.4 (d, 1H), 3.5 (s, 3H), 3.65 (d, 1H), 3.9 (d, 1H), 6.65 (d, 1H), 6.75 (d, 1H), 7.15 (dd, 1H), 7.3 (m, 5H).

10 Anal. calc'd for C<sub>24</sub>H<sub>30</sub>N<sub>4</sub>OS•3HCl: C, 54.19; H, 6.25; N, 10.53. Found: C, 53.91; H, 6.39; N, 10.27.

EXAMPLE 55

{[4-Methoxy-3-(2-phenylpiperidin-3-ylaminomethyl)  
phenyl]-methyl-sulfamoyl}-acetic acid ethyl ester

15 48% yield, m.p. 245-248°C.

MS: m/e 475 (M<sup>+</sup>, 5%) 356, 175, 150 (100%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, free base) δ 1.3 (t, 3H), 1.3502.15 (m, 6H), 2.8 (m, 2H), 3.3 (d, 1H), 3.35 (s, 3H), 3.4 (d, 1H), 3.5 (s, 3H), 3.65 (d, 1H), 3.9 (d, 3H), 4.3 (q, 2H), 6.7 (d, 20 1H), 7.15 (d, 1H), 7.35 (m, 6H).

EXAMPLE 56

2-Hydroxyethanesulfonic acid [4-methoxy-3-(2-  
phenylpiperidin-3-ylaminomethyl)phenyl]-methanamide  
hydrochloride

25 4% yield, m.p. 255-260°C (dec.).

MS: m/e (433, M<sup>+</sup>), 314 (85%), 258 (100%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>, free base) δ 2.55 (bs, 4H), 2.75 (t, 1H), 2.85 (m, 1H), 3.15 (t, 2H), 3.2 (s, 3H), 3.35 (d, 1H), 3.5 (s, 3H), 3.65 (d, 1H), 3.9 (d, 1H), 3.95 (t, 2H), 6.65 (d, 30 1H), 7.1-7.4 (m, 7H).

Anal. calc'd for C<sub>22</sub>H<sub>31</sub>N<sub>3</sub>O<sub>4</sub>S•HCl: C, 52.17; H, 6.57; N, 8.29. Found: C 51.89, N 6.27, N 7.95.

EXAMPLE 57

N-(3,4-Dichlorobenzyl)-N-[4-methoxy-3-(2-  
phenylpiperidin-3-ylaminomethyl)phenyl]-methanesulfonamide  
dihydrochloride hydrate

35 13% yield, m.p. 240-243°C (dec.).



-70-

MS: m/e 548 ( $M^{+1}$ , 8%), 428 (30), 159 (100).

$^1H$  NMR ( $CDCl_3$ , free base)  $\delta$  1.35-2.15 (m, 6H), 2.65 (m, 1H), 2.8 (t, 1H), 2.85 (s, 3H), 3.25 (d, 1H), 3.35 (d, 1H), 3.5 (s, 3H), 3.65 (d, 1H), 3.9 (d, 1H), 4.65 (q, 2H), 6.6 (d, 1H), 6.9 (d, 1H), 7.0 (dd, 1H), 7.15 (dd, 1H), 7.2-7.4 (m, 7H).

Anal. calc'd for  $C_{27}H_{31}Cl_2N_3O_3S \cdot 2HCl \cdot 2/3H_2O$ : C, 51.19; H, 5.46; N, 6.63. Found: C, 51.17; H, 5.33; N, 6.48.

#### EXAMPLE 58

10 4,5-Dimethylthiazole-2-sulfonic acid methyl-[3-(2-phenylpiperidin-3-yl-aminomethyl)-4-trifluoromethoxyphenyl]-amide trihydrochloride hydrate

12% yield, m.p. 239-240°C (dec.),

MS: m/e 555 ( $M^{+1}$ ), 380.

15  $^1H$  NMR ( $CDCl_3$ , free base)  $\delta$  1.5 (m, 1H), 1.7 (m, 1H), 1.9 (m, 4H), 2.1 (m, 1H), 2.2 (s, 3H), 2.7 (s, 3H), 2.8 (m, 2H), 3.2 (s, 3H), 3.3 (m, 1H), 3.5 (q, 2H), 3.9 (d, 1H), 7.0 (m, 3H), 7.2 (m, 5H).

Anal. calc'd for  $C_{25}H_{29}F_3N_4O_3S_2 \cdot 3HCl \cdot H_2O$ : C, 44.09; H, 4.88; N, 8.23. Found: C, 44.36; H, 4.95; N, 8.51.

The title compounds of examples 59-62 were prepared by a procedure similar to that of Example 38C, starting with the appropriate aldehyde in place of 2-methoxy-5-methylthiobenzaldehyde.

25

#### EXAMPLE 59

(2S,3S)-3-[2-Methoxy-5-(N-acethyl-N-methylamino)benzyl-aminol]-2-diphenylmethyl-1-azabicyclo[2.2.2]octane dihydrochloride

M.p.: 232-234°C (AcOEt).

30 IR(KBr): 3430, 3055, 3020, 1648, 1500, 1386, 1244, 709  $cm^{-1}$ .

$^1H$  NMR (270 MHz,  $CDCl_3$ , free amine): 7.36-7.07 (m, 10H), 6.95 (dd, J=8.6, 2.6 Hz, 1H), 6.71 (d, J=8.6 Hz, 1H), 6.37 (d, J=2.6 Hz, 1H), 4.49 (d, J=12.1 Hz, 1H), 3.78-3.71 (m, 1H), 3.65-3.60 (m, 1H), 3.63 (s, 3H), 3.28-3.23 (m, 2H), 3.20 (s, 3H), 2.93 (dd, J=7.7, 4.4 Hz, 1H), 2.81 (m, 2H), 2.68 (m, 1H), 2.04 (m, 1H), 1.82 (s, 3H), 1.95-1.29 (m, 5H).

-71-

EXAMPLE 60

(2S,3S)-3-[2-Methoxy-5-(N-methyl-N-trifluoro-acetyl-amino)benzylamino]-2-diphenylmethyl-1-azabicyclo[2.2.2]octane dihydrochloride

5       <sup>1</sup>H NMR (270 MHz, CDCl<sub>3</sub>, ppm) (free amine): 7.38-7.04 (m, 10H), 7.01 (dd, J=2.6, 8.8 Hz, 1H), 6.69 (d, J=8.8 Hz, 1H), 6.42 (br.s, 1H), 4.48 (d, J=12.1 Hz, 1H), 3.77-3.55 (m, 2H), 3.61 (s, 3H), 3.44-3.15 (m, 2H), 3.28 (s, 3H), 2.89 (dd, J=4.0, 7.7 Hz, 1H), 2.86-2.60 (m, 3H), 2.05-1.82 (m, 10 2H), 1.75-1.40 (m, 2H), 1.38-1.20 (m, 1H).

IR (KBr, cm<sup>-1</sup>) (free amine): 3360, 1699, 1598, 1499, 1465, 1451, 1248, 1203, 1150, 1112, 1071, 1038, 817, 754, 703.

EXAMPLE 61

15       (2S,3S)-3-[5-(N-Isopropyl)-N-methylsulfonylamino]-2-methoxybenzylamino]-2-diphenylmethyl-1-azabicyclo[2.2.2]octane dihydrochloride

M.p.: 178-179°C.

<sup>1</sup>H NMR (270 MHz, CDCl<sub>3</sub>, ppm) (free amine): 7.34-7.03 (m, 20 10H), 7.07 (dd, J=2.6, 8.8 Hz, 1H), 6.82 (d, J=2.6 Hz, 1H), 6.68 (d, J=8.8 Hz, 1H), 4.53 (sep, J=6.6 Hz, 1H), 4.49 (d, J=12.1 Hz, 1H), 3.76-3.63 (m, 1H), 3.62 (d, J=13.6 Hz, 1H), 3.53 (s, 3H), 3.30-3.13 (m, 1H), 3.23 (d, J=13.6 Hz, 1H), 2.97-2.86 (m, 1H), 2.93 (s, 3H), 2.84-2.58 (m, 3H), 2.11-25 2.02 (m, 1H), 2.00-1.40 (m, 4H), 1.38-1.20 (m, 1H), 1.15 (dd, J=2.6, 6.6 Hz, 6H).

IR (KBr, cm<sup>-1</sup>) (free amine): 3340, 1603, 1495, 1462, 1450, 1366, 1332, 1232, 1181, 1154, 1130, 1107, 1032, 982, 961, 815, 801, 755, 703.

30

EXAMPLE 62

(2S,3S)-3-[2-Methoxy-5-(N-methyl-N-methylsulfonylamino)benzylamino]-2-diphenylmethyl-1-azabicyclo[2.2.2]octane monomethanesulfonate

M.p.: 197-203°C (IPA-Hex).

35       IR (KBr): 3430, 2945, 1500, 1340, 1218, 1166, 1039, 746 cm<sup>-1</sup>.


-72-

<sup>1</sup>H NMR (270 MHz, CDCl<sub>3</sub>, free amine): 7.35-7.07 (m, 11H), 6.82 (d, J=2.6 Hz, 1H), 6.68 (d, J=8.8 Hz, 1H), 4.49 (d, J=12.1 Hz, 1H), 3.76-3.67 (m, 1H), 3.61-3.53 (m, 1H), 3.54 (s, 3H), 3.26 (s, 3H), 3.26-3.18 (m, 2H), 2.93 (dd, 5 J=7.7, 4.0 Hz, 1H), 2.83 (s, 3H), 2.82-2.77 (m, 2H), 2.65 (m, 1H), 2.06 (m, 1H), 1.91-1.55 (m, 4H), 1.34-1.23 (m, 1H).

-73-

CLAIMS

1. A compound of the formula



wherein ring A is an aryl group selected from phenyl, naphthyl, thienyl, dihydroquinoliny and indoliny, and  
 10 wherein the side chain containing  $\text{NR}^2\text{R}^3$  is attached to a carbon atom of ring A;

W is hydrogen,  $(\text{C}_1\text{-C}_6)$ alkyl,  $\text{S-}(\text{C}_1\text{-C}_3)$ alkyl, halo or  $(\text{C}_1\text{-C}_6)$ alkoxy optionally substituted with from one to three fluorine atoms;

15  $\text{R}^1$  is selected from amino,  $(\text{C}_1\text{-C}_6)$ alkylamino, di- $(\text{C}_1\text{-C}_6)$ alkylamino,  $\text{-S(O)}_v\text{-(C}_1\text{-C}_{10})$ alkyl wherein v is zero, one or two,  $\text{-S(O)}_v\text{-aryl}$  wherein v is zero, one or two,  $\text{-O-aryl}$ ,  $\text{-SO}_2\text{NR}^4\text{R}^5$  wherein each of  $\text{R}^4$  and  $\text{R}^5$  is, independently,  $(\text{C}_1\text{-C}_6)$ alkyl, or  $\text{R}^4$  and  $\text{R}^5$ , together with the nitrogen to  
 20 which they are attached, form a saturated ring containing one nitrogen and from 3 to 6

25 carbons,  $\text{-NHC(=O)(C}_1\text{-C}_6\text{)alkyl}$ ,  $\text{-N[(C}_1\text{-C}_6\text{)alkyl-C(=O)(C}_1\text{-C}_6\text{)alkyl}$ ,


$\text{-NHCCF}_3$ ,  $\text{-N[(C}_1\text{-C}_6\text{)alkyl-CCF}_3$ ,  $(\text{C}_1\text{-C}_{10})\text{alkyl-N-SO}_2\text{-(C}_1\text{-C}_{10})\text{alkyl}$   
 30 wherein one or both of the alkyl moieties may optionally be substituted with from one to three fluorine atoms,

$\text{-N(SO}_2\text{-(C}_1\text{-C}_{10})\text{alkyl)}_2$  and  $(\text{C}_1\text{-C}_{10})\text{alkyl-N-SO}_2\text{-aryl}$ ; and wherein  
 35 the aryl moieties of said  $\text{-S(O)}_v\text{-aryl}$ ,  $\text{-O-aryl}$  and

$(\text{C}_1\text{-C}_{10})\text{alkyl-N-SO}_2\text{-aryl}$  are independently selected from phenyl and benzyl and may optionally be substituted with  
 40 from one to three substituents independently selected from  $(\text{C}_1\text{-C}_4)$ alkyl,  $(\text{C}_1\text{-C}_4)$ alkoxy and halo;

-74-

or R<sup>1</sup> is a group having the formula





wherein a is 0, 1 or 2 and the asterisk represents a position meta to the R<sup>2</sup>R<sup>3</sup>NCH<sub>2</sub> side chain;

R<sup>2</sup> is hydrogen or -CO<sub>2</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl;



10 R<sup>3</sup> is selected from

-75-


5




10




15




20



25



30



35

-76-


wherein  $R^6$  and  $R^{10}$  are independently selected from furyl, thienyl, pyridyl, indolyl, biphenyl and phenyl, wherein said phenyl may optionally be substituted with one or two substituents independently selected from halo,  $(C_1-C_{10})$  alkyl optionally substituted with from one to three fluorine atoms,  $(C_1-C_{10})$  alkoxy optionally substituted with from one to three fluorine atoms, carboxy, benzyloxycarbonyl and  $(C_1-C_3)$  alkoxy-carbonyl;

$R^7$  is selected from  $(C_3-C_4)$  branched alkyl,  $(C_5-C_6)$  branched alkenyl,  $(C_5-C_7)$  cycloalkyl, and the radicals named in the definition of  $R^6$ ;

$R^8$  is hydrogen or  $(C_1-C_6)$  alkyl;

$R^9$  and  $R^{19}$  are independently selected from phenyl, biphenyl, naphthyl, pyridyl, benzhydryl, thienyl or furyl, and  $R^9$  and  $R^{19}$  may optionally be substituted with from one to three substituents independently selected from halo,  $(C_1-C_{10})$  alkyl optionally substituted with from one to three fluorine atoms and  $(C_1-C_{10})$  alkoxy optionally substituted with from one to three fluorine atoms;

$Y$  is  $(CH_2)_1$ , wherein 1 is an integer from one to three, or  $Y$  is a group of the formula



$Z$  is oxygen, sulfur, amino,  $(C_1-C_3)$  alkylamino or  $(CH_2)_n$ , wherein  $n$  is zero, one or two;

$x$  is zero, one or two;  
 $y$  is zero, one or two;  
 $z$  is three, four or five;  
 $o$  is two or three;  
 $p$  is zero or one;  
 $r$  is one, two or three;

-77-

the ring containing  $(CH_2)_1$ , may contain from zero to three double bonds, and one of the carbon atoms of  $(CH_2)_2$ , may optionally be replaced by oxygen, sulfur or nitrogen;

$R^{11}$  is thienyl, biphenyl or phenyl optionally substituted with one or two substituents independently selected from halo,  $(C_1-C_{10})$  alkyl optionally substituted with from one to three fluorine atoms and  $(C_1-C_{10})$  alkoxy optionally substituted with from one to three fluorine atoms;

10 X is  $(CH_2)_q$ , wherein q is an integer from 1 to 6, and wherein any one of the carbon-carbon single bonds in said  $(CH_2)_q$ , may optionally be replaced by a carbon-carbon double bond, and wherein any one of the carbon atoms of said  $(CH_2)_q$ , may optionally be substituted with  $R^{14}$ , and wherein any one  
15 of the carbon atoms of said  $(CH_2)_q$ , may optionally be substituted with  $R^{15}$ ;





m is an integer from 0 to 8, and any one of the carbon-carbon single bonds of  $(CH_2)_m$ , wherein both carbon atoms of such bond are bonded to each other and to another  
20 carbon atom in the  $(CH_2)_m$  chain, may optionally be replaced by a carbon-carbon double bond or a carbon-carbon triple bond, and any one of the carbon atoms of said  $(CH_2)_m$  may optionally be substituted with  $R^{17}$ ;

$R^{12}$  is a radical selected from hydrogen,  $(C_1-C_6)$  straight or branched alkyl,  $(C_3-C_7)$ cycloalkyl wherein one of the carbon atoms may optionally be replaced by nitrogen, oxygen or sulfur; aryl selected from biphenyl, phenyl, indanyl and naphthyl; heteroaryl selected from thienyl, furyl, pyridyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, triazolyl, tetrazolyl and quinolyl; phenyl- $(C_2-C_6)$ alkyl, benzhydryl and benzyl, wherein the point of attachment on  $R^{12}$  is a carbon atom unless  $R^{12}$  is hydrogen, and wherein each of said aryl and heteroaryl groups and the phenyl moieties of said benzyl, phenyl- $(C_2-C_6)$ alkyl and benzhydryl may optionally be  
35 substituted with one or more substituents independently selected from halo, nitro,  $(C_1-C_{10})$ alkyl optionally



-78-



substituted with from one to three fluorine atoms,  
 (C<sub>1</sub>-C<sub>10</sub>)alkoxy optionally substituted with  
 from one to three fluorine atoms, amino, hydroxy-(C<sub>1</sub>-  
 C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)-alkylamino,



- R<sup>13</sup> is hydrogen, phenyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl;


or R<sup>12</sup> and R<sup>13</sup>, together with the carbon to which they  
 are attached, form a saturated carbocyclic ring having from  
 30 3 to 7 carbon atoms wherein one of said carbon atoms that is  
 neither the point of attachment of the spiro ring nor  
 adjacent to it may optionally be replaced by oxygen,  
 nitrogen or sulfur;

R<sup>14</sup> and R<sup>15</sup> are each independently selected from  
 35 hydrogen, hydroxy, halo, amino, oxo (=O), cyano, hydroxy-(C<sub>1</sub>-  
 C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino,




45

-79-



5

10 set forth in the definition of  $R^{12}$ ;

$R^{16}$  is  $NH\overset{\overset{O}{\parallel}}{C}R^{18}$ ,  $NHCH_2R^{18}$ ,  $SO_2R^{18}$ ,  $CO_2H$  or one of the radicals  
15 set forth in any of the definitions of  $R^{12}$ ,  $R^{14}$  and  $R^{15}$ ;

$R^{17}$  is oximino ( $=NOH$ ) or one of the radicals set forth  
in any of the definitions of  $R^{12}$ ,  $R^{14}$  and  $R^{15}$ ; and

$R^{18}$  is  $(C_1-C_6)\text{alkyl}$ , hydrogen, phenyl or phenyl  $(C_1-C_6)\text{alkyl}$ ;

20 with the proviso that (a) when  $m$  is 0, one of  $R^{16}$  and  $R^{17}$   
is absent and the other is hydrogen, (b) when  $R^3$  is a group  
of the formula VIII,  $R^{14}$  and  $R^{15}$  cannot be attached to the  
same carbon atom, (c) when  $R^{14}$  and  $R^{15}$  are attached to the  
same carbon atom, then either each of  $R^{14}$  and  $R^{15}$  is  
25 independently selected from hydrogen, fluoro,  $(C_1-C_6)\text{alkyl}$ ,  
hydroxy- $(C_1-C_6)\text{alkyl}$  and  $(C_1-C_6)\text{alkoxy}-(C_1-C_6)\text{alkyl}$ , or  $R^{14}$  and  
 $R^{15}$ , together with the carbon to which they are attached,  
form a  $(C_3-C_6)$  saturated carbocyclic ring that forms a spiro  
compound with the nitrogen-containing ring to which they are  
30 attached; (d) when  $R^1$  is amino,  $(C_1-C_6)\text{alkylamino}$ , di-

$(C_1-C_6)\text{alkylamino}$  or  $-NH\overset{\overset{O}{\parallel}}{C}(C_1-C_6)\text{alkyl}$ ,  $R^3$  is a group of the  
35 formula II, III, IV, V or VI; and (e) when  $R^{14}$  or  $R^{15}$  is  
attached to a carbon atom of X or  $(CH_2)$ , that is adjacent to  
the ring nitrogen, then  $R^{14}$  or  $R^{15}$ , respectively, must be a  
substituent wherein the point of attachment is a carbon  
atom;

40 or a pharmaceutically acceptable salt of such compound.

2. A compound according to claim 1, wherein  $R^3$  is a  
group of the formula II, III, VII or IX;  $R^2$  is hydrogen; ring

-80-

A is phenyl or indolinyl; W is (C<sub>1</sub>-C<sub>3</sub>)alkoxy optionally substituted with from one to three fluorine atoms; and R<sup>1</sup> is S(O)<sub>v</sub>-(C<sub>1</sub>-C<sub>10</sub>)alkyl wherein v is zero, one or two, S(O)<sub>v</sub>-aryl

5

wherein v is zero, one or two, -O-aryl, (C<sub>1</sub>-C<sub>10</sub>)alkyl-N-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>10</sub>)alkyl wherein one or both of the alkyl moieties may optionally be substituted with from one to three fluorine

10

atoms, -N(SO<sub>2</sub>-C<sub>1</sub>-C<sub>10</sub>)alkyl)<sub>2</sub> or (C<sub>1</sub>-C<sub>10</sub>)alkyl-N-SO<sub>2</sub>-aryl

15 wherein said aryl is phenyl or benzyl and may optionally be substituted with from one to three substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy and halo.

3. A compound according to claim 2, wherein R<sup>3</sup> is a group of the formula II, o is two, and each R<sup>6</sup> and R<sup>7</sup> is phenyl.

4. A compound according to claim 2, wherein R<sup>3</sup> is a group of the formula VII, each of R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup> and R<sup>16</sup> is hydrogen, m is zero and X is -(CH<sub>2</sub>)<sub>3</sub>-.

25 5. A compound according to claim 2, wherein R<sup>3</sup> is a group of the formula IX, R<sup>19</sup> is benzhydryl and r is two.

6. A compound according to claim 2, wherein R<sup>3</sup> is a group of the formula III, R<sup>8</sup> is other than hydrogen and R<sup>9</sup> is benzyhydryl.

30 7. A compound according to claim 1, wherein the substituents at positions "2" and "3" of the nitrogen containing ring are in the cis configuration.

8. A compound according to claim 1, wherein R<sup>3</sup> is a group of the formula II wherein the substituents at positions "2" and "3" of the nitrogen containing ring are in the cis configuration, o is two, each of R<sup>6</sup> and R<sup>7</sup> is phenyl and ring A is phenyl or indolinyl.

9. A compound according to claim 1, wherein R<sup>3</sup> is a group of the formula III wherein the substituents at positions "2" and "3" of the nitrogen containing ring are in

40

-81-

the cis configuration,  $R^8$  is other than hydrogen,  $R^9$  is benzhydryl and ring A is phenyl.

10. A compound according to claim 1, wherein  $R^3$  is a group of the formula VII wherein  $R^{12}$  and the substituent at position "3" of the nitrogen containing ring are in the cis configuration, ring A is phenyl,  $R^{12}$  is phenyl, each of  $R^2$ ,  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$  and  $R^{16}$  is hydrogen, m is zero, X is  $-(CH_2)_2-$  or  $-(CH_2)_3-$  and  $R^1$  is selected from  $S(O)_v-(C_1-C_{10})$ alkyl wherein v is zero, one or two, and  $(C_1-C_{10})$ alkyl-N-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>10</sub>)alkyl, and di-(C<sub>1</sub>-C<sub>6</sub>)alkylamino.

11. A compound according to claim 10, wherein X is  $-(CH_2)_2-$  and W is (C<sub>1</sub>-C<sub>6</sub>) alkoxy optionally substituted with from one to three fluorine atoms.

12. A compound according to claim 10, wherein X is  $-(CH_2)_3-$  and W is (C<sub>1</sub>-C<sub>6</sub>) alkoxy optionally substituted with from one to three fluorine atoms.

13. A compound according to claim 1 wherein  $R^3$  is a group of the formula IX wherein the substituents at positions "2" and "3" of the nitrogen containing ring are in the cis configuration, r is two and  $R^{19}$  is benzhydryl.

14. A compound according to claim 13, wherein ring A is phenyl, W is (C<sub>1</sub>-C<sub>6</sub>) alkoxy optionally substituted with from one to three fluorine atoms and  $R^1$  is selected from  $-S(O)_v-(C_1-C_{10})$ alkyl wherein v is zero, one or two, di-(C<sub>1</sub>-C<sub>6</sub>)alkylamino and  $(C_1-C_{10})$ alkyl-N-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>10</sub>)alkyl.


15. A compound according to claim 6, wherein ring A is phenyl, W is (C<sub>1</sub>-C<sub>6</sub>)alkoxy optionally substituted with from one to three fluorine atoms, and  $R^1$  is selected from  $-S(O)_v-(C_1-C_{10})$ alkyl wherein v is zero, one or two, and  $(C_1-C_{10})$ alkyl-N-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>10</sub>)alkyl.

16. A compound according to claim 6, wherein ring A is phenyl, W is (C<sub>1</sub>-C<sub>6</sub>)alkoxy optionally substituted with from

-82-

one to three fluorine atoms, and  $R^2$  is selected from amino,  $(C_1-C_6)$ alkylamino or di- $(C_1-C_6)$ alkylamino.

17. A compound according to claim 3, wherein ring A is phenyl, W is  $(C_1-C_6)$ alkoxy optionally substituted with from one to three fluorine atoms, and  $R^1$  is selected from  $-S(O)_v-$   $(C_1-C_{10})$ alkyl wherein v is zero, one or two, and  $(C_1-C_{10})$ alkyl-

 N-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>10</sub>)alkyl.

10 18. A compound according to claim 3, wherein ring A is phenyl, W is  $(C_1-C_6)$ alkoxy optionally substituted with from one to three fluorine atoms, and  $R^2$  is selected from amino,  $(C_1-C_6)$ alkylamino or di- $(C_1-C_6)$ alkylamino.

19. A compound according to claim 15, wherein W is attached at the "2" position of ring A and  $R^1$  is attached at the "5" position of ring A, relative to the point of attachment of the  $NR^2R^3$  containing side chain.

20. A compound according to claim 16, wherein W is attached at the "2" position of ring A and  $R^1$  is attached at the "5" position of ring A, relative to the point of attachment of the  $NR^2R^3$  containing side chain.

21. A compound according to claim 17, wherein W is attached at the "2" position of ring A and  $R^1$  is attached at the "5" position of ring A, relative to the point of attachment of the  $NR^2R^3$  containing side chain.

22. A compound according to claim 18, wherein W is attached at the "2" position of ring A and  $R^1$  is attached at the "5" position of ring A, relative to the point of attachment of the  $NR^2R^3$  containing side chain.

23. A compound according to claim 15, wherein W is selected from isopropoxy,  $OCF_3$ ,  $OCH_3$ ,  $OCHF_2$  and  $OCH_2CF_3$ .

24. A compound according to claim 16, wherein W is selected from isopropoxy,  $OCF_3$ ,  $OCH_3$ ,  $OCHF_2$  and  $OCH_2CF_3$ .

25. A compound according to claim 17, wherein W is selected from isopropoxy,  $OCF_3$ ,  $OCH_3$ ,  $OCHF_2$  and  $OCH_2CF_3$ .

26. A compound according to claim 18, wherein W is selected from isopropoxy,  $OCF_3$ ,  $OCH_3$ ,  $OCHF_2$  and  $OCH_2CF_3$ .

-83-

27. A compound according to claim 4, wherein ring A is phenyl, W is selected from isopropoxy,  $\text{OCF}_3$ ,  $\text{OCH}_3$ ,  $\text{OCHF}_2$  and  $\text{OCH}_2\text{CF}_3$ , and  $\text{R}^1$  is selected from  $-\text{S}(\text{O})_v-(\text{C}_1-\text{C}_{10})\text{alkyl}$  wherein v

5 is zero, one or two, and  $(\text{C}_1-\text{C}_{10})\text{alkyl}-\overset{\text{vw}}{\text{N}}-\text{SO}_2-(\text{C}_1-\text{C}_{10})\text{alkyl}$ .

28. (2S,3S)-N-(2-methoxy-5-methylsulfonylphenyl)-methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine.

29. (2S,3S)-N-(2-methoxy-5-methylthiophenyl)methyl-2-  
10 diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine.


30. (2S,3S)-N-(2-methoxy-5-dimethylaminophenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine.

31. (2S,3S)-N-(5-trifluoroacetyl-amino-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo-  
15 [2.2.2]octan-3-amine.

32. A compound according to claim 1, wherein  $\text{R}^3$  is a group of the formula VII, m is zero, each of  $\text{R}^{13}$ ,  $\text{R}^{15}$ ,  $\text{R}^{16}$  and

20  $\text{R}^{17}$  is hydrogen,  $\text{R}^{12}$  is phenyl,  $\text{R}^{14}$  is  $-\overset{\text{O}}{\parallel}\text{C}-\text{OH}$ , ring A is phenyl, W is  $(\text{C}_1-\text{C}_3)\text{alkoxy}$  and  $\text{R}^1$  is selected from  $(\text{C}_1-\text{C}_3)\text{alkyl}$ ,  $-\text{SCH}_3$ ,  $\text{SO}_2\text{CH}_3$ ,  $\text{SOCH}_3$ ,  $(\text{C}_1-\text{C}_6)\text{alkylamino}$  and di- $(\text{C}_1-\text{C}_6)\text{alkyl-amino}$ .

33. A compound according to claim 32, wherein said  
25 compound has the formula



35 34. A compound according to claim 1, wherein said compound is selected from the group consisting of:

-84-

- (2S, 3S) - 3 - [2-methoxy-5-(N-acetyl-N-methylamino)benzylamino]-2-diphenylmethyl-1-azabicyclo[2.2.2]octane;
- (2S,3S)-N-(5-ethylthio-2-methoxyphenyl)methyl-2-  
5 diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(5-ethylsulfinyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(5-ethylsulfonyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- 10 (2S,3S)-N-(5-isopropylthio-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(5-isopropylsulfinyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(5-isopropylsulfonyl-2-methoxyphenyl)methyl-  
15 2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(5-diethylamino-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S, 3S) - N - [2-methoxy-5-(N-methyl-N-methanesulfonylamino)phenyl]methyl-2-diphenylmethyl-1-  
20 azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-[5-N-isopropylsulfonyl-N-methylamino)-2-methoxyphenyl]methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-[5-(N-isopropyl-N-methanesulfonylamino)-2-  
25 methoxyphenyl]methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (2S,3S)-N-(5-amino-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine;
- (1SR,2SR,3SR,4RS)-3-(2-methoxy-5-(N-methyl-N-methanesulfonylamino)benzyl)amino-2-benzhydryl-[2.2.1]-  
30 azanorbornane;
- (1SR,2SR,3SR,4RS)-3-(2-isopropoxy-5-(N-methyl-N-methanesulfonyl-amino)benzyl)amino-2-benzhydryl-[2.2.1]-azanorbornane;
- 35 (1SR,2SR,3SR,4RS)-3-(2-methoxy-5-(N-methyl-N-trifluoromethane-sulfonylamino)benzyl)amino-2-benzhydryl-[2.2.1]-azanorbornane;

-85-

(1SR, 2SR, 3SR, 4RS) -3-(2-methoxy-5-(N-methyl-N-phenylmethane-sulfonylamino)benzyl)amino-2-benzhydryl-[2.2.1]-azanorbornane;

(1SR, 2SR, 3SR, 4RS) -3-(2-methoxy-5-(N-isopropyl-N-methane-sulfonylamino)benzyl)amino-2-benzhydryl-[2.2.1]-azanorbornane;

(1SR, 2SR, 3SR, 4RS) -3-(2-methoxy-5-(1,1-dioxo-2-isothiazolidinyl)benzyl)amino-2-benzhydryl-[2.2.1]-azanorbornane;

10 (1SR, 2SR, 3SR, 4RS) -3-[(2,3-dihydro-5-methoxy-1-methanesulfonyl-6-indolyl)methylamino]-2-benzhydryl-[2.2.1]-azanorbornane;

cis-3-(2-methylthiobenzyl)amino-2-phenylpiperidine;

cis-3-(5-chloro-2-methylthiobenzyl)amino-2-phenylpiperidine;

cis-3-(5-fluoro-2-methylthiobenzyl)amino-2-phenylpiperidine;

cis-3-(5-tert-butyl-2-methylthiobenzyl)amino-2-phenylpiperidine;

20 cis-3-(2-tert-butylthiobenzyl)amino-2-phenylpiperidine;

cis-3-(2-methylsulfonylbenzyl)amino-2-phenylpiperidine;

cis-3-(2-methoxy-5-methylthiobenzyl)amino-2-phenylpiperidine;

cis-3-(2-difluoromethoxy-5-methylthiobenzyl)amino-2-phenylpiperidine;

cis-3-(2-methoxy-5-methylsulfinylbenzyl)amino-2-phenylpiperidine;

cis-3-(2-methoxy-5-methylsulfonylbenzyl)amino-2-phenylpiperidine;

30 cis-3-[2-methoxy-5-(N,N-diethylaminosulfonyl)benzyl]amino-2-phenylpiperidine;

cis-3-[2-isopropoxy-5-(N,N-diethylaminosulfonyl)benzyl]amino-2-phenylpiperidine;

cis-3-(2-(4-chlorophenylthio)benzyl)amino-2-phenylpiperidine;

cis-3-(2-phenyloxybenzylamino)-2-phenylpiperidine;



-86-

- cis-3-(2-methoxy-5-phenyloxybenzylamino)-2-phenylpiperidine;
- cis-3-[2-methoxy-5-(N-methyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- 5 cis-3-[2-trifluoromethoxy-5-(N-methyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- cis-3-[2-isopropoxy-5-(N-methyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- cis-3-[2-cyclopentyloxy-5-(N-methyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- 10 cis-3-[2-methoxy-5-(N-isopropyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- cis-3-[2-isopropoxy-5-(N-isopropyl-N-methanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- 15 cis-3-[2-methoxy-5-(N-cyclopentyl-N-methanesulfonylamino)amino-2-phenylpiperidine;
- cis-3-[2-methoxy-5-(N-methyl-N-trifluoromethanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- cis-3-[2-isopropoxy-5-(N-methyl-N-trifluoromethanesulfonylamino)benzyl]amino-2-phenylpiperidine;
- 20 cis-3-[2-methoxy-5-(N-methyl-N-isopropylsulfonylamino)benzyl]amino-2-phenylpiperidine;
- cis-3-[2-methoxy-5-(N-methyl-N-(4-methylphenylsulfonyl)amino)benzyl]amino-2-phenylpiperidine;
- 25 cis-3-[2-isopropoxy-5-(N-methyl-N-(4-methylphenylsulfonyl)amino)benzyl]amino-2-phenylpiperidine;
- cis-3-[2-methoxy-5-(N-methyl-N-phenylmethylsulfonylamino)benzyl]amino-2-phenylpiperidine;
- cis-3-[2-trifluoromethoxy-5-(N,N-bis(methanesulfonyl)amino)benzyl]amino-2-phenylpiperidine;
- 30 cis-3-[2-methoxy-5-(1,1-dioxo-2-isothiazolidinyl)benzyl]amino-2-phenyl-piperidine;
- cis-3-[(2,3-dihydro-5-methoxy-1-methanesulfonyl-6-indolyl)methylamino]-2-phenylpiperidine;
- 35 cis-3-[(2,3-dihydro-5-methoxy-2-methyl-1-methanesulfonyl-6-indolyl)methylamino]-2-phenylpiperidine;

-87-


(2SR,3SR,4RS)-2-benzhydryl-4-(2-hydroxyethyl)-3-(2-methoxy-5-methylthiobenzyl)aminopyrrolidine;

(2SR,3SR,4RS)-2-benzhydryl-4-(2-hydroxyethyl)-3-(2-methoxy-5-(N-methyl-N-methanesulfonylamino)benzyl)amino-  
5 pyrrolidine;


(2SR,3SR,4RS)-2-benzhydryl-4-(2-hydroxyethyl)-3-(2-methoxy-5-(N-thiazolidine-S,S-dioxide)benzyl)aminopyrrolidine and;

cis-3-(2-ethylthiophenyl)methylamino-2-phenyl-  
10 piperidine.

35. A compound of the formula



or



25 wherein  $R^1$ ,  $W$ , ring  $A$  and  $R^3$  are defined as in claim 1.

36. A pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases, anxiety, colitis, depression or dysthymic disorders, psychosis, pain, allergies, chronic  
30 obstructive airways disease, hypersensitivity disorders, vasospastic diseases, fibrosing and collagen diseases, reflex sympathetic dystrophy, addiction disorders, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders, disorders related to immune  
35 enhancement or suppression and rheumatic diseases in a mammal, comprising an amount of a compound according to

-88-

claim 1 effective in preventing or treating such condition and a pharmaceutically acceptable carrier.

37. A method of treating or preventing a condition selected from the group consisting of inflammatory diseases  
5 anxiety, colitis, depression or dysthymic disorders, psychosis, pain, allergies, chronic obstructive airways disease, hypersensitivity disorders, vasospastic diseases, fibrosing and collagen diseases, reflex sympathetic dystrophy, addiction disorders, stress related somatic  
10 disorders, peripheral neuropathy, neuralgia, neuropathological disorders, disorders related to immune enhancement or suppression and rheumatic diseases in a mammal, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to  
15 claim 1 effective in preventing or treating such condition.

38. A pharmaceutical composition for antagonizing the substance P receptor in a mammal, comprising a substance P receptor antagonizing effective amount of a compound according to claim 1 and a pharmaceutically acceptable  
20 carrier.

39. A method of antagonizing the substance P receptor in a mammal, comprising administering to said mammal a substance P receptor antagonizing effective amount of a compound according to claim 1.

25 40. A pharmaceutical composition for treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of a compound according to claim 1 effective in  
30 antagonizing the effect of substance P at its receptor site and a pharmaceutically acceptable carrier.

41. A method of treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated  
35 neurotransmission, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1, or a pharmaceutically acceptable salt

-89-

thereof, effective in antagonizing the effect of substance P at its receptor site.

42. A pharmaceutical composition for treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such condition and a pharmaceutically acceptable carrier.

43. A method of treating or preventing a condition in mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1 effective in treating or preventing such condition.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 93/04063

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5	C07D211/56; C07D453/02;	C07D417/12; A61K31/445; C07D401/12; A61K31/435; C07D487/08 /(C07D453/02, ..
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
Int.Cl. 5	C07D ; A61K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
Y	WO,A,9 201 688 (PFIZER INC.) 6 February 1992 cited in the application *SEE WHOLE APPLICATION, ESPECIALLY DEFINITION OF R1*	1-36, 38, 40, 42
Y	WO,A,9 118 899 (PFIZER INC.) 12 December 1991 cited in the application *SEE WHOLE APPLICATION, ESPECIALLY DEFINITION OF R2*	1-36, 38, 40, 42
P,X	WO,A,9 300 331 (PFIZER INC.) 7 January 1993 cited in the application *SEE DEFINITIONS OF Q, AND X1-X3*	1, 10-14, 35-36, 38, 40, 42
-/--		
<p><sup>10</sup> Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&amp;" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
30 AUGUST 1993		14. 09. 93
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		SCRUTON-EVANS I.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 93/04063

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup> According to International Patent Classification (IPC) or to both National Classification and IPC IPC <sup>5</sup> : 221:00, 211:00) (C07D487/08, 209:00, 209:00)								
<b>II. FIELDS SEARCHED</b> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black; margin: 5px 0;">Minimum Documentation Searched <sup>7</sup></div> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%; border: 1px solid black; padding: 5px;">Classification System</td> <td style="border: 1px solid black; padding: 5px;">Classification Symbols</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px; height: 40px; vertical-align: top;">IPC<sup>5</sup></td> <td style="border: 1px solid black;"></td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black; margin: 5px 0;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup></div>			Classification System	Classification Symbols	IPC <sup>5</sup>			
Classification System	Classification Symbols							
IPC <sup>5</sup>								
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 10%; border: 1px solid black; padding: 5px;">Category <sup>10</sup></th> <th style="width: 60%; border: 1px solid black; padding: 5px;">Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup></th> <th style="width: 30%; border: 1px solid black; padding: 5px;">Relevant to Claim No. <sup>13</sup></th> </tr> <tr> <td style="border: 1px solid black; height: 300px;"></td> <td style="border: 1px solid black;"></td> <td style="border: 1px solid black;"></td> </tr> </table>			Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>			
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>						
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p><sup>10</sup> Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Δ" document member of the same patent family</p> </div> </div>								
<b>IV. CERTIFICATION</b> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border: 1px solid black; padding: 5px;">Date of the Actual Completion of the International Search</td> <td style="width: 50%; border: 1px solid black; padding: 5px;">Date of Mailing of this International Search Report</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;">           International Searching Authority  <div style="text-align: center; margin-top: 10px;">EUROPEAN PATENT OFFICE</div> </td> <td style="border: 1px solid black; padding: 5px;">           Signature of Authorized Officer  <div style="text-align: center; margin-top: 10px;"> </div> </td> </tr> </table>			Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	International Searching Authority <div style="text-align: center; margin-top: 10px;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: center; margin-top: 10px;"> </div>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report							
International Searching Authority <div style="text-align: center; margin-top: 10px;">EUROPEAN PATENT OFFICE</div>	Signature of Authorized Officer <div style="text-align: center; margin-top: 10px;"> </div>							

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
Y	WO,A,9 206 079 (PFIZER INC.) 16 April 1992 cited in the application *SEE DEFINITIONS OF R3* ---	1-36,38, 40,42
P,Y	WO,A,9 301 170 (PFIZER INC.) 21 January 1993 *SEE DEFINITIONS OF R3* ---	1-36,38, 40,42
P,X	WO,A,9 220 676 (PFIZER INC.) 26 November 1992 cited in the application *SEE DEFINITIONS OF Ar1 AND W* ---	1,2,6,9, 36,38, 40,42
P,X	WO,A,9 217 449 (PFIZER INC.) 15 October 1992 *SEE DEFINITIONS OF R1* -----	1,2,4, 10,12

**ANNEX TO THE INTERNATIONAL SEARCH REPORT  
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9304063  
SA 74083

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

30/08/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9201688	06-02-92	AU-A- 7894891 CN-A- 1058405 EP-A- 0540526	18-02-92 05-02-92 12-05-93
WO-A-9118899	12-12-91	AU-A- 7860291 EP-A- 0532527	31-12-91 24-03-93
WO-A-9300331	07-01-93	AU-A- 1889392 CN-A- 1067655	25-01-93 06-01-93
WO-A-9206079	16-04-92	AU-A- 8746391 CA-A- 2089736 CN-A- 1060285 EP-A- 0550635	28-04-92 29-03-92 15-04-92 14-07-93
WO-A-9301170	21-01-93	US-A- 5232929	03-08-93
WO-A-9220676	26-11-92	AU-A- 1927592	30-12-92
WO-A-9217449	15-10-92	AU-A- 1244892 CN-A- 1065264	02-11-92 14-10-92